



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C07K 16/00, A61K 39/395, C12N 15/13, 15/85, 5/10, G01N 33/68, 33/577 // C07K 16/28	A2	(11) International Publication Number: WO 00/42072 (43) International Publication Date: 20 July 2000 (20.07.00)
(21) International Application Number: PCT/US00/00973 (22) International Filing Date: 14 January 2000 (14.01.00) (30) Priority Data: 60/116,023 15 January 1999 (15.01.99) US (71) Applicant: GENENTECH, INC. [US/US]; 1 DNA Way, South San Francisco, CA 94080-4990 (US). (72) Inventor: PRESTA, Leonard, G.; 1900 Gough Street, #206, San Francisco, CA 94109 (US). (74) Agents: LEE, Wendy, M. et al.; Genentech, Inc., 1 DNA Way, South San Francisco, CA 94080-4990 (US).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: POLYPEPTIDE VARIANTS WITH ALTERED EFFECTOR FUNCTION (57) Abstract The present invention concerns polypeptides comprising a variant Fc region. More particularly, the present invention concerns Fc region-containing polypeptides that have altered effector function as a consequence of one or more amino acid modifications in the Fc region thereof.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Larvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

POLYPEPTIDE VARIANTS WITH ALTERED EFFECTOR FUNCTION**BACKGROUND OF THE INVENTION****Field of the Invention**

5 The present invention concerns polypeptides comprising a variant Fc region. More particularly, the present invention concerns Fc region-containing polypeptides that have altered effector function as a consequence of one or more amino acid modifications in the Fc region thereof.

Description of Related Art

10 Antibodies are proteins which exhibit binding specificity to a specific antigen. Native antibodies are usually heterotetrameric glycoproteins of about 150,000 daltons, composed of two identical light (L) chains and two identical heavy (H) chains. Each light chain is linked to a heavy chain by one covalent disulfide bond, while the number of disulfide linkages varies between the heavy chains of different immunoglobulin isotypes. Each heavy and light chain also has regularly spaced intrachain disulfide bridges. Each heavy chain has at one end a variable domain (V_H)
15 followed by a number of constant domains. Each light chain has a variable domain at one end (V_L) and a constant domain at its other end; the constant domain of the light chain is aligned with the first constant domain of the heavy chain, and the light chain variable domain is aligned with the variable domain of the heavy chain. Particular amino acid residues are believed to form an interface between the light and heavy chain variable domains.

20 The term "variable" refers to the fact that certain portions of the variable domains differ extensively in sequence among antibodies and are responsible for the binding specificity of each particular antibody for its particular antigen. However, the variability is not evenly distributed through the variable domains of antibodies. It is concentrated in three segments called complementarity determining regions (CDRs) both in the light chain and the heavy chain variable
25 domains. The more highly conserved portions of the variable domains are called the framework regions (FRs). The variable domains of native heavy and light chains each comprise four FRs, largely adopting a β -sheet configuration, connected by three CDRs, which form loops connecting, and in some cases forming part of, the β -sheet structure. The CDRs in each chain are held together in close proximity by the FRs and, with the CDRs from the other chain, contribute to the
30 formation of the antigen binding site of antibodies (see Kabat *et al.*, *Sequences of Proteins of Immunological Interest*, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, MD. (1991)).

The constant domains are not involved directly in binding an antibody to an antigen, but exhibit various effector functions. Depending on the amino acid sequence of the constant region
35 of their heavy chains, antibodies or immunoglobulins can be assigned to different classes. There

are five major classes of immunoglobulins: IgA, IgD, IgE, IgG and IgM, and several of these may be further divided into subclasses (isotypes), e.g. IgG1, IgG2, IgG3, and IgG4; IgA1 and IgA2.

The heavy chain constant regions that correspond to the different classes of immunoglobulins are called α , δ , ϵ , γ , and μ , respectively. Of the various human immunoglobulin classes, only human IgG1, IgG2, IgG3 and IgM are known to activate complement; and human IgG1 and IgG3 mediate ADCC more effectively than IgG2 and IgG4.

A schematic representation of the native IgG1 structure is shown in Fig. 1, where the various portions of the native antibody molecule are indicated. Papain digestion of antibodies produces two identical antigen binding fragments, called Fab fragments, each with a single antigen binding site, and a residual "Fc" fragment, whose name reflects its ability to crystallize readily. The crystal structure of the human IgG Fc region has been determined (Deisenhofer, *Biochemistry* 20:2361-2370 (1981)). In human IgG molecules, the Fc region is generated by papain cleavage N-terminal to Cys 226. The Fc region is central to the effector functions of antibodies.

The effector functions mediated by the antibody Fc region can be divided into two categories: (1) effector functions that operate after the binding of antibody to an antigen (these functions involve the participation of the complement cascade or Fc receptor (FcR)-bearing cells); and (2) effector functions that operate independently of antigen binding (these functions confer persistence in the circulation and the ability to be transferred across cellular barriers by transcytosis). Ward and Ghetie, *Therapeutic Immunology* 2:77-94 (1995).

While binding of an antibody to the requisite antigen has a neutralizing effect that might prevent the binding of a foreign antigen to its endogenous target (e.g. receptor or ligand), binding alone may not remove the foreign antigen. To be efficient in removing and/or destructing foreign antigens, an antibody should be endowed with both high affinity binding to its antigen, and efficient effector functions.

Fc receptor (FcR) binding

The interaction of antibodies and antibody-antigen complexes with cells of the immune system effects a variety of responses, including antibody-dependent cell-mediated cytotoxicity (ADCC) and complement dependent cytotoxicity (CDC) (reviewed in Daëron, *Annu. Rev. Immunol.* 15:203-234 (1997); Ward and Ghetie, *Therapeutic Immunol.* 2:77-94 (1995); as well as Ravetch and Kinet, *Annu. Rev. Immunol.* 9:457-492 (1991)).

Several antibody effector functions are mediated by Fc receptors (FcRs), which bind the Fc region of an antibody. FcRs are defined by their specificity for immunoglobulin isotypes; Fc receptors for IgG antibodies are referred to as Fc γ R, for IgE as Fc ϵ R, for IgA as Fc α R and so on. Three subclasses of Fc γ R have been identified: Fc γ RI (CD64), Fc γ RII (CD32) and Fc γ RIII (CD16).

Because each FcγR subclass is encoded by two or three genes, and alternative RNA splicing leads to multiple transcripts, a broad diversity in FcγR isoforms exists. The three genes encoding the FcγRI subclass (FcγRIA, FcγRIB and FcγRIC) are clustered in region 1q21.1 of the long arm of chromosome 1; the genes encoding FcγRII isoforms (FcγRIIA, FcγRIIB and FcγRIIC) and the two genes encoding FcγRIII (FcγRIIIA and FcγRIIIB) are all clustered in region 1q22. These different FcR subtypes are expressed on different cell types (reviewed in Ravetch and Kinet, *Annu. Rev. Immunol.* 9:457-492 (1991)). For example, in humans, FcγRIIIB is found only on neutrophils, whereas FcγRIIIA is found on macrophages, monocytes, natural killer (NK) cells, and a subpopulation of T-cells. Notably, FcγRIIIA is the only FcR present on NK cells, one of the cell types implicated in ADCC.

FcγRI, FcγRII and FcγRIII are immunoglobulin superfamily (IgSF) receptors; FcγRI has three IgSF domains in its extracellular domain, while FcγRII and FcγRIII have only two IgSF domains in their extracellular domains.

Another type of Fc receptor is the neonatal Fc receptor (FcRn). FcRn is structurally similar to major histocompatibility complex (MHC) and consists of an α-chain noncovalently bound to β2-microglobulin.

The binding site on human and murine antibodies for FcγR have been previously mapped to the so-called "lower hinge region" consisting of residues 233-239 (EU index numbering as in Kabat *et al.*, *Sequences of Proteins of Immunological Interest*, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, MD. (1991)). Woof *et al. Molec. Immunol.* 23:319-330 (1986); Duncan *et al. Nature* 332:563 (1988); Canfield and Morrison, *J. Exp. Med.* 173:1483-1491 (1991); Chappel *et al., Proc. Natl. Acad. Sci USA* 88:9036-9040 (1991). Of residues 233-239, P238 and S239 have been cited as possibly being involved in binding, but these two residues have never been evaluated by substitution or deletion.

Other previously cited areas possibly involved in binding to FcγR are: G316-K338 (human IgG) for human FcγRI (by sequence comparison only; no substitution mutants were evaluated) (Woof *et al. Molec. Immunol.* 23:319-330 (1986)); K274-R301 (human IgG1) for human FcγRIII (based on peptides) (Sarmay *et al. Molec. Immunol.* 21:43-51 (1984)); Y407-R416 (human IgG) for human FcγRIII (based on peptides) (Gergely *et al. Biochem. Soc. Trans.* 12:739-743 (1984)); as well as N297 and E318 (murine IgG2b) for murine FcγRII (Lund *et al., Molec. Immunol.*, 29:53-59 (1992)).

Pro331 in IgG3 was changed to Ser, and the affinity of this variant to target cells analyzed. The affinity was found to be six-fold lower than that of unmutated IgG3, indicating the involvement of Pro331 in FcγRI binding. Morrison *et al., Immunologist*, 2:119-124 (1994); and Canfield and Morrison, *J. Exp. Med.* 173:1483-91 (1991).

C1q binding

C1q and two serine proteases, C1r and C1s, form the complex C1, the first component of the complement dependent cytotoxicity (CDC) pathway. C1q is a hexavalent molecule with a molecular weight of approximately 460,000 and a structure likened to a bouquet of tulips in which six collagenous "stalks" are connected to six globular head regions. Burton and Woof, *Advances in Immunol.* 51:1-84 (1992). To activate the complement cascade, it is necessary for C1q to bind to at least two molecules of IgG1, IgG2, or IgG3 (the consensus is that IgG4 does not activate complement), but only one molecule of IgM, attached to the antigenic target. Ward and Ghetie, *Therapeutic Immunology* 2:77-94 (1995) at page 80.

Based upon the results of chemical modifications and crystallographic studies, Burton *et al.* (*Nature*, 288:338-344 (1980)) proposed that the binding site for the complement subcomponent C1q on IgG involves the last two (C-terminal) β -strands of the CH2 domain. Burton later suggested (*Molec. Immunol.*, 22(3):161-206 (1985)) that the region comprising amino acid residues 318 to 337 might be involved in complement fixation.

Duncan and Winter (*Nature* 332:738-40 (1988)), using site directed mutagenesis, reported that Glu318, Lys320 and Lys322 form the binding site to C1q. The data of Duncan and Winter were generated by testing the binding of a mouse IgG2b isotype to guinea pig C1q. The role of Glu318, Lys320 and Lys322 residues in the binding of C1q was confirmed by the ability of a short synthetic peptide containing these residues to inhibit complement mediated lysis. Similar results are disclosed in U.S. Patent No. 5,648,260 issued on July 15, 1997, and U.S. Patent No. 5,624,821 issued on April 29, 1997.

The residue Pro331 has been implicated in C1q binding by analysis of the ability of human IgG subclasses to carry out complement mediated cell lysis. Mutation of Ser331 to Pro331 in IgG4 conferred the ability to activate complement. (Tao *et al.*, *J. Exp. Med.*, 178:661-667 (1993); Brekke *et al.*, *Eur. J. Immunol.*, 24:2542-47 (1994)).

From the comparison of the data of the Winter group, and the Tao *et al.* and Brekke *et al.* papers, Ward and Ghetie concluded in their review article that there are at least two different regions involved in the binding of C1q: one on the β -strand of the CH2 domain bearing the Glu318, Lys320 and Lys322 residues, and the other on a turn located in close proximity to the same β -strand, and containing a key amino acid residue at position 331.

Other reports suggested that human IgG1 residues Leu235, and Gly237, located in the lower hinge region, play a critical role in complement fixation and activation. Xu *et al.*, *Immunol.* 150:152A (Abstract) (1993). WO94/29351 published December 22, 1994 reports that amino acid residues necessary for C1q and FcR binding of human IgG1 are located in the N-terminal region of the CH2 domain, *i.e.* residues 231 to 238.

It has further been proposed that the ability of IgG to bind C1q and activate the complement cascade also depends on the presence, absence, or modification of the carbohydrate moiety positioned between the two CH2 domains (which is normally anchored at Asn297). Ward and Ghetie, *Therapeutic Immunology* 2:77-94 (1995) at page 81.

5

SUMMARY OF THE INVENTION

The present invention provides a variant of a parent polypeptide comprising an Fc region, which variant mediates antibody-dependent cell-mediated cytotoxicity (ADCC) in the presence of human effector cells more effectively, or binds an Fc gamma receptor (FcγR) with better affinity, than the parent polypeptide and comprises at least one amino acid modification in the Fc region.

10 The polypeptide variant may, for example, comprise an antibody or an immunoadhesin. The Fc region of the parent polypeptide preferably comprises a human Fc region; *e.g.*, a human IgG1, IgG2, IgG3 or IgG4 Fc region. The polypeptide variant preferably comprises an amino acid modification (*e.g.* a substitution) at any one or more of amino acid positions 256, 290, 298, 312, 326, 330, 333, 334, 360, 378 or 430 of the Fc region, wherein the numbering of the residues in
15 the Fc region is that of the EU index as in Kabat.

In addition, the invention provides a polypeptide comprising a variant Fc region with altered Fc gamma receptor (FcγR) binding affinity, which polypeptide comprises an amino acid modification at any one or more of amino acid positions 238, 239, 248, 249, 252, 254, 255, 256, 258, 265, 267, 268, 269, 270, 272, 276, 278, 280, 283, 285, 286, 289, 290, 292, 293, 294, 295,
20 296, 298, 301, 303, 305, 307, 309, 312, 315, 320, 322, 324, 326, 327, 329, 330, 331, 333, 334, 335, 337, 338, 340, 360, 373, 376, 378, 382, 388, 389, 398, 414, 416, 419, 430, 434, 435, 437, 438 or 439 of the Fc region, wherein the numbering of the residues in the Fc region is that of the EU index as in Kabat. The variant Fc region preferably comprises a variant human IgG Fc region, *e.g.*, a variant human IgG1, IgG2, IgG3 or IgG4 Fc region. In this respect, it is noted that, in the
25 work in the above-cited art where the parent polypeptide had a non-human murine Fc region, different residues from those identified herein were thought to impact FcR binding. For example, in the murine IgG2b/murine FcγRII system, IgG E318 was found to be important for binding (Lund *et al. Molec. Immunol.* 27(1):53-59 (1992)), whereas E318A had no effect in the human IgG/human FcγRII system (Table 6 below).

30 In one embodiment, the polypeptide variant with altered FcγR binding activity displays reduced binding to an FcγR and comprises an amino acid modification at any one or more of amino acid positions 238, 239, 248, 249, 252, 254, 265, 268, 269, 270, 272, 278, 289, 292, 293, 294, 295, 296, 298, 301, 303, 322, 324, 327, 329, 333, 335, 338, 340, 373, 376, 382, 388, 389, 414, 416, 419, 434, 435, 437, 438 or 439 of the Fc region, wherein the numbering of the residues
35 in the Fc region is that of the EU index as in Kabat.

For example, the polypeptide variant may display reduced binding to an FcγRI and comprise an amino acid modification at any one or more of amino acid positions 238, 265, 269, 270, 327 or 329 of the Fc region, wherein the numbering of the residues in the Fc region is that of the EU index as in Kabat.

5 The polypeptide variant may display reduced binding to an FcγRII and comprise an amino acid modification at any one or more of amino acid positions 238, 265, 269, 270, 292, 294, 295, 298, 303, 324, 327, 329, 333, 335, 338, 373, 376, 414, 416, 419, 435, 438 or 439 of the Fc region, wherein the numbering of the residues in the Fc region is that of the EU index as in Kabat.

10 The polypeptide variant of interest may display reduced binding to an FcγRIII and comprise an amino acid modification at one or more of amino acid positions 238, 239, 248, 249, 252, 254, 265, 268, 269, 270, 272, 278, 289, 293, 294, 295, 296, 301, 303, 322, 327, 329, 338, 340, 373, 376, 382, 388, 389, 416, 434, 435 or 437 of the Fc region, wherein the numbering of the residues in the Fc region is that of the EU index as in Kabat.

15 In another embodiment, the polypeptide variant with altered FcγR binding affinity displays improved binding to the FcγR and comprises an amino acid modification at any one or more of amino acid positions 255, 256, 258, 267, 268, 272, 276, 280, 283, 285, 286, 290, 298, 301, 305, 307, 309, 312, 315, 320, 322, 326, 330, 331, 333, 334, 337, 340, 360, 378, 398 or 430 of the Fc region, wherein the numbering of the residues in the Fc region is that of the EU index as in Kabat.

20 For example, the polypeptide variant may display increased binding to an FcγRIII and, optionally, may further display decreased binding to an FcγRII. An exemplary such variant comprises amino acid modification(s) at position(s) 298 and/or 333 of the Fc region, wherein the numbering of the residues in the Fc region is that of the EU index as in Kabat.

25 The polypeptide variant may display increased binding to an FcγRII and comprise an amino acid modification at any one or more of amino acid positions 255, 256, 258, 267, 268, 272, 276, 280, 283, 285, 286, 290, 301, 305, 307, 309, 312, 315, 320, 322, 326, 330, 331, 337, 340, 378, 398 or 430 of the Fc region, wherein the numbering of the residues in the Fc region is that of the EU index as in Kabat. Such polypeptide variants with increased binding to an FcγRII may optionally further display decreased binding to an FcγRIII and may, for example, comprise an amino acid modification at any one or more of amino acid positions 268, 272, 298, 301, 322 or 30 340 of the Fc region, wherein the numbering of the residues in the Fc region is that of the EU index as in Kabat.

35 The invention further provides a polypeptide comprising a variant Fc region with altered neonatal Fc receptor (FcRn) binding affinity, which polypeptide comprises an amino acid modification at any one or more of amino acid positions 238, 252, 253, 254, 255, 256, 265, 272, 286, 288, 303, 305, 307, 309, 311, 312, 317, 340, 356, 360, 362, 376, 378, 380, 382, 386, 388,

400, 413, 415, 424, 433, 434, 435, 436, 439 or 447 of the Fc region, wherein the numbering of the residues in the Fc region is that of the EU index as in Kabat. Such polypeptide variants with reduced binding to an FcRn may comprise an amino acid modification at any one or more of amino acid positions 252, 253, 254, 255, 288, 309, 386, 388, 400, 415, 433, 435, 436, 439 or 447
5 of the Fc region, wherein the numbering of the residues in the Fc region is that of the EU index as in Kabat. The above-mentioned polypeptide variants may, alternatively, display increased binding to FcRn and comprise an amino acid modification at any one or more of amino acid positions 238, 256, 265, 272, 286, 303, 305, 307, 311, 312, 317, 340, 356, 360, 362, 376, 378, 380, 382, 413, 424 or 434 of the Fc region, wherein the numbering of the residues in the Fc region
10 is that of the EU index as in Kabat.

The invention also provides a composition comprising the polypeptide variant and a physiologically or pharmaceutically acceptable carrier or diluent. This composition for potential therapeutic use is sterile and may be lyophilized.

Diagnostic and therapeutic uses for the polypeptide variants disclosed herein are
15 contemplated. In one diagnostic application, the invention provides a method for determining the presence of an antigen of interest comprising exposing a sample suspected of containing the antigen to the polypeptide variant and determining binding of the polypeptide variant to the sample. In one therapeutic application, the invention provides a method of treating a mammal suffering from or predisposed to a disease or disorder, comprising administering to the mammal
20 a therapeutically effective amount of a polypeptide variant as disclosed herein, or of a composition comprising the polypeptide variant and a pharmaceutically acceptable carrier.

The invention further provides: isolated nucleic acid encoding the polypeptide variant; a vector comprising the nucleic acid, optionally, operably linked to control sequences recognized by a host cell transformed with the vector; a host cell containing the vector; a method for
25 producing the polypeptide variant comprising culturing this host cell so that the nucleic acid is expressed and, optionally, recovering the polypeptide variant from the host cell culture (e.g. from the host cell culture medium).

The invention further provides a method for making a variant Fc region with altered Fc receptor (FcR) binding affinity, or altered antibody-dependent cell-mediated cytotoxicity (ADCC)
30 activity, comprising:

- (a) introducing one or more amino acid modifications into an Fc region of a parent polypeptide in order to generate a variant Fc region;
- (b) determining binding of the variant Fc region to an FcR, or determining ADCC activity of the variant Fc region.

Step (b) of the method may comprise determining binding of the variant Fc region to one or more FcRs *in vitro*. Moreover, the method may result in the identification of a variant Fc region with improved FcR binding affinity, or with improved ADCC activity, in step (b) thereof. Where step (b) comprises determining binding of the Fc region to an FcR, the FcR may, for example, be human Fc gamma receptor III (FcγRIII). Where step (b) comprises determining binding of the variant Fc region to at least two different FcRs, the FcRs tested preferably include human Fc gamma receptor II (FcγRII) and human Fc gamma receptor III (FcγRIII).

Brief Description of the Drawings

Figure 1 is a schematic representation of a native IgG. Disulfide bonds are represented by heavy lines between CH1 and CL domains and the two CH2 domains. V is variable domain; C is constant domain; L stands for light chain and H stands for heavy chain.

Figure 2 shows C1q binding of wild type (wt) C2B8 antibody; C2B8 antibody with a human IgG2 constant region (IgG2); and variants K322A, K320A and E318A.

Figure 3 depicts C1q binding of variants P331A, P329A and K322A.

Figures 4A and 4B depict the amino acid sequences of E27 anti-IgE antibody light chain (Fig. 4A; SEQ ID NO:1) and heavy chain (Fig. 4B; SEQ ID NO:2).

Figure 5 is a schematic diagram of the "immune complex" prepared for use in the FcR assay described in Example 1. The hexamer comprising three anti-IgE antibody molecules (the "Fc region-containing polypeptide") and three IgE molecules (the "first target molecule") is shown. IgE has two "binding sites" for the anti-IgE antibody (E27) in the Fc region thereof. Each IgE molecule in the complex is further able to bind two VEGF molecules ("the second target polypeptide"). VEGF has two "binding sites" for IgE.

Figure 6 shows C1q binding results obtained for variants D270K and D270V compared to wild type C2B8.

Figure 7 depicts complement dependent cytotoxicity (CDC) of variants D270K and D270V, compared to wild type C2B8.

Figure 8 shows C1q binding ELISA results for 293 cell-produced wild type C2B8 antibody (293-Wt-C2B8), CHO-produced wild type C2B8 antibody (CHO-Wt-C2B8) and various variant antibodies.

Figure 9 shows C1q binding ELISA results obtained for wild type (wt) C2B8 and various variant antibodies as determined in Example 3.

Figure 10 depicts the three-dimensional structure of a human IgG Fc region, highlighting residues: Asp270, Lys326, Pro329, Pro331, Lys322 and Glu333.

Figure 11 shows C1q binding ELISA results obtained for wild type C2B8 and various variant antibodies as determined in Example 3.

Figure 12 shows C1q binding ELISA results obtained for wild type C2B8 and double variants, K326M-E333S and K326A-E333A.

Figure 13 shows CDC of wild type C2B8 and double variants, K326M-E333S and K326A-E333A.

5 Figure 14 depicts C1q binding ELISA results obtained for C2B8 with a human IgG4 (IgG4), wild type C2B8 (Wt-C2B8), C2B8 with a human IgG2 constant region (IgG2), and variant antibodies as described in Example 3.

10 Figures 15A and 15B show binding patterns for parent antibody (E27) to Fc γ RIIB and Fc γ RIIIA. Figure 15A shows the binding pattern for the humanized anti-IgE E27 IgG1 as a monomer (open circles), hexamer (closed squares), and immune complex consisting of multiple hexamers (closed triangles) to a recombinant GST fusion protein of the human Fc γ RIIB (CD32) receptor α subunit. The hexameric complex (closed squares) was formed by the mixture of equal molar concentrations of E27 (which binds to the Fc region of human IgE) and a human myeloma IgE. The hexamer is a stable 1.1 kD complex consisting of 3 IgG molecules (150 kD each) and
15 3 IgE molecules (200 kD each). The immune complex (closed triangles) was formed sequentially by first mixing equal molar concentrations of E27 and recombinant anti-VEGF IgE (human IgE with Fab variable domains that bind human VEGF) to form the hexamer. Hexamers were then linked to form an immune complex by the addition of 2x molar concentration of human VEGF, a 44 kD homodimer which has two binding sites for the anti-VEGF IgE per mole of VEGF. Figure
20 15B shows the binding pattern to a recombinant GST fusion protein of the human Fc γ RIIIA (CD16) receptor α subunit.

Figure 16A shows the binding of immune complexes using different antigen-antibody pairs to recombinant GST fusion protein of the Fc γ RIIA receptor α subunit. Figure 16B shows the binding of the same antigen-antibody pairs to the GST fusion protein of the Fc γ RIIIA receptor α subunit. Closed circles represent binding of human IgE:anti-IgE E27 IgG1; open circles represent
25 binding of human VEGF:humanized anti-VEGF IgG1.

Figure 17 summarizes differences in binding selectivity of some alanine variants between the different Fc γ Rs. Binding of alanine variants at residues in the CH2 domain of anti-IgE E27 IgG1 are shown to Fc γ RIIA, Fc γ RIIB, and Fc γ RIIIA. Type 1 abrogates binding to all three
30 receptors: D278A (265 in EU numbering). Type 2 improves binding to Fc γ RIIA and Fc γ RIIB, while binding to Fc γ RIIIA is unaffected: S280A (267 in EU numbering). Type 3 improves binding to Fc γ RIIA and Fc γ RIIB, but reduces binding to Fc γ RIIIA: H281A (268 in EU numbering). Type 4 reduces binding to Fc γ RIIA and Fc γ RIIB, while improving binding to Fc γ RIIIA: S317A (298 in EU numbering). Type 5 improves binding to Fc γ RIIIA, but does not affect binding to Fc γ RIIA and
35 Fc γ RIIB: E352A, K353A (333 and 334 in EU numbering).

Figures 18A and 18B compare the FcγRIIIA protein/protein assay and CHO GPI-FcγRIIIA cell based assay, respectively. Figure 18A illustrates binding of selected alanine variants to FcγRIIIA-GST fusion protein. S317A (298 in EU numbering) and S317A/K353A (298 and 334 in EU numbering) bind better than E27 wildtype, while D278A (265 in EU numbering) almost
 5 completely abrogates binding. Figure 18B illustrates that a similar pattern of binding is found on CHO cells expressing a recombinant GPI-linked form of FcγRIIIA.

Figures 19A and 19B compare the FcγRIIB protein/protein assay and CHO GPI-FcγRIIB cell based assay, respectively. Figure 19A illustrates binding of selected alanine variants to FcγRIIB-GST fusion protein. H281A (268 in EU numbering) binds better than E27 wildtype while
 10 S317A (298 in EU numbering) shows reduced binding. Figure 19B illustrates that a similar pattern of binding is found on CHO cells expressing a recombinant membrane bound form of FcγRIIB.

Figure 20 shows single alanine substitutions in the CH2 domain of anti-HER2 IgG1 (HERCEPTIN®) that influence FcγRIIIA binding in both the protein-protein and cell-based assays alter the ability to bind to FcγRIIIA on peripheral blood mononuclear cell (PBMC) effector cells.
 15 Recombinant humanized anti-HER2 (HERCEPTIN®), which binds to HER2-expressing SK-BR-3 breast tumor cells, was preincubated with ⁵¹Cr-labeled SK-BR-3 cells for 30 minutes (opsonization) at 100 ng/ml (filled circles) and 1.25 ng/ml (filled squares). Keeping the SK-BR-3 tumor target cell concentration constant, the ratio of effector cells was increased from 0 to 100. The spontaneous cytotoxicity in the absence of antibody (hatched squares) was 20% at an
 20 effector:target (E:T) ratio of 100:1. A single alanine mutation that did not affect FcγRIIIA binding, variant G31 = R309A (292 in EU numbering), did not effect ADCC (filled triangles). A single alanine mutation that only slightly increased binding to FcγRIIIA, variant G30 = K307A (290 in EU numbering), also showed slightly improved ADCC (*i.e.*, a 1.1 fold improvement in ADCC activity, calculated as area under the curve) at 1.25 ng/ml at all E:T ratios (filled diamonds) compared to
 25 wildtype antibody at 1.25 ng/ml (filled square). A single alanine mutation that decreased binding to FcγRIIIA, variant G34 = Q312A (295 in EU numbering), also showed decreased ADCC activity (filled inverted triangles).

Figure 21 illustrates that a single alanine mutation which had the most improved binding to FcγRIIIA, variant G36 = S317A (298 in EU numbering), in the protein-protein and cell-based
 30 assays also showed the most improvement in ADCC (filled triangles) among the variants compared to wildtype (closed squares) at 1.25 ng/ml. G36 displayed a 1.7 fold improvement in ADCC activity, calculated as area under the curve. Variants G17 = E282A (269 in EU numbering) and G18 = D283A (270 in EU numbering) both showed reduced binding to FcγRIIIA as well as reduced efficacy in ADCC. The effector cells were PBMCs.

Figure 22A depicts alignments of native sequence IgG Fc regions. Native sequence human IgG Fc region sequences, humIgG1 (non-A and A allotypes) (SEQ ID NOs: 3 and 4, respectively), humIgG2 (SEQ ID NO:5), humIgG3 (SEQ ID NO:6) and humIgG4 (SEQ ID NO:7), are shown. The human IgG1 sequence is the non-A allotype, and differences between this sequence and the A allotype (at positions 356 and 358; EU numbering system) are shown below the human IgG1 sequence. Native sequence murine IgG Fc region sequences, murIgG1 (SEQ ID NO:8), murIgG2A (SEQ ID NO:9), murIgG2B (SEQ ID NO:10) and murIgG3 (SEQ ID NO:11), are also shown. Figure 22B shows percent identity among the Fc region sequences of Figure 22A.

Figure 23 depicts alignments of native sequence human IgG Fc region sequences, humIgG1 (non-A and A allotypes; SEQ ID NOs:3 and 4, respectively), humIgG2 (SEQ ID NO:5), humIgG3 (SEQ ID NO:6) and humIgG4 (SEQ ID NO:7) with differences between the sequences marked with asterisks.

Figure 24 shows area under curve (AUC) for selected variants compared to anti-HER2 IgG1 (HERCEPTIN®) in a 4 hour ADCC assay. The effector cells were PBMCs (N=5). Variant G36 (S317A; 298 in Eu numbering) with improved binding to FcγRIIIA showed improved ADCC activity; variant G31 (R309A; 292 in Eu numbering) which did not display altered FcγRIIIA binding, also had unaltered ADCC activity; and G14 (D265A; 278 in Eu numbering) which had reduced FcγRIIIA binding, also had reduced ADCC activity.

Detailed Description of the Preferred Embodiments

I. Definitions

Throughout the present specification and claims, the numbering of the residues in an immunoglobulin heavy chain is that of the EU index as in Kabat *et al.*, *Sequences of Proteins of Immunological Interest*, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, MD (1991), expressly incorporated herein by reference. The "EU index as in Kabat" refers to the residue numbering of the human IgG1 EU antibody.

A "parent polypeptide" is a polypeptide comprising an amino acid sequence which lacks one or more of the Fc region modifications disclosed herein and which differs in effector function compared to a polypeptide variant as herein disclosed. The parent polypeptide may comprise a native sequence Fc region or an Fc region with pre-existing amino acid sequence modifications (such as additions, deletions and/or substitutions).

The term "Fc region" is used to define a C-terminal region of an immunoglobulin heavy chain, *e.g.*, as shown in Figure 1. The "Fc region" may be a native sequence Fc region or a variant Fc region. Although the boundaries of the Fc region of an immunoglobulin heavy chain might vary, the human IgG heavy chain Fc region is usually defined to stretch from an amino acid

residue at position Cys226, or from Pro230, to the carboxyl-terminus thereof. The Fc region of an immunoglobulin generally comprises two constant domains, CH2 and CH3, as shown, for example, in Fig. 1.

5 The "CH2 domain" of a human IgG Fc region (also referred to as "C_γ2" domain) usually extends from about amino acid 231 to about amino acid 340. The CH2 domain is unique in that it is not closely paired with another domain. Rather, two N-linked branched carbohydrate chains are interposed between the two CH2 domains of an intact native IgG molecule. It has been speculated that the carbohydrate may provide a substitute for the domain-domain pairing and help stabilize the CH2 domain. Burton, *Molec. Immunol.*22:161-206 (1985).

10 The "CH3 domain" comprises the stretch of residues C-terminal to a CH2 domain in an Fc region (*i.e.* from about amino acid residue 341 to about amino acid residue 447 of an IgG)

A "functional Fc region" possesses an "effector function" of a native sequence Fc region. Exemplary "effector functions" include C1q binding; complement dependent cytotoxicity; Fc receptor binding; antibody-dependent cell-mediated cytotoxicity (ADCC); phagocytosis; down
15 regulation of cell surface receptors (*e.g.* B cell receptor; BCR), etc. Such effector functions generally require the Fc region to be combined with a binding domain (*e.g.* an antibody variable domain) and can be assessed using various assays as herein disclosed, for example.

A "native sequence Fc region" comprises an amino acid sequence identical to the amino acid sequence of an Fc region found in nature. Native sequence human Fc regions are shown
20 in Fig. 23 and include a native sequence human IgG1 Fc region (non-A and A allotypes); native sequence human IgG2 Fc region; native sequence human IgG3 Fc region; and native sequence human IgG4 Fc region as well as naturally occurring variants thereof. Native sequence murine Fc regions are shown in Fig. 22A.

A "variant Fc region" comprises an amino acid sequence which differs from that of a native
25 sequence Fc region by virtue of at least one "amino acid modification" as herein defined. Preferably, the variant Fc region has at least one amino acid substitution compared to a native sequence Fc region or to the Fc region of a parent polypeptide, *e.g.* from about one to about ten amino acid substitutions, and preferably from about one to about five amino acid substitutions in a native sequence Fc region or in the Fc region of the parent polypeptide. The variant Fc region
30 herein will preferably possess at least about 80% homology with a native sequence Fc region and/or with an Fc region of a parent polypeptide, and most preferably at least about 90% homology therewith, more preferably at least about 95% homology therewith.

"Homology" is defined as the percentage of residues in the amino acid sequence variant that are identical after aligning the sequences and introducing gaps, if necessary, to achieve the
35 maximum percent homology. Methods and computer programs for the alignment are well known

in the art. One such computer program is "Align 2", authored by Genentech, Inc., which was filed with user documentation in the United States Copyright Office, Washington, DC 20559, on December 10, 1991.

5 The term "Fc region-containing polypeptide" refers to a polypeptide, such as an antibody or immunoadhesin (see definitions below), which comprises an Fc region.

The terms "Fc receptor" or "FcR" are used to describe a receptor that binds to the Fc region of an antibody. The preferred FcR is a native sequence human FcR. Moreover, a preferred FcR is one which binds an IgG antibody (a gamma receptor) and includes receptors of the FcγRI, FcγRII, and FcγRIII subclasses, including allelic variants and alternatively spliced forms
10 of these receptors. FcγRII receptors include FcγRIIA (an "activating receptor") and FcγRIIB (an "inhibiting receptor"), which have similar amino acid sequences that differ primarily in the cytoplasmic domains thereof. Activating receptor FcγRIIA contains an immunoreceptor tyrosine-based activation motif (ITAM) in its cytoplasmic domain. Inhibiting receptor FcγRIIB contains an immunoreceptor tyrosine-based inhibition motif (ITIM) in its cytoplasmic domain. (see review M.
15 in Daëron, *Annu. Rev. Immunol.* 15:203-234 (1997)). FcRs are reviewed in Ravetch and Kinet, *Annu. Rev. Immunol* 9:457-92 (1991); Capel *et al.*, *Immunomethods* 4:25-34 (1994); and de Haas *et al.*, *J. Lab. Clin. Med.* 126:330-41 (1995). Other FcRs, including those to be identified in the future, are encompassed by the term "FcR" herein. The term also includes the neonatal receptor, FcRn, which is responsible for the transfer of maternal IgGs to the fetus (Guyer *et al.*,
20 *J. Immunol.* 117:587 (1976) and Kim *et al.*, *J. Immunol.* 24:249 (1994)).

"Antibody-dependent cell-mediated cytotoxicity" and "ADCC" refer to a cell-mediated reaction in which nonspecific cytotoxic cells that express FcRs (*e.g.* Natural Killer (NK) cells, neutrophils, and macrophages) recognize bound antibody on a target cell and subsequently cause lysis of the target cell. The primary cells for mediating ADCC, NK cells, express FcγRIII
25 only, whereas monocytes express FcγRI, FcγRII and FcγRIII. FcR expression on hematopoietic cells is summarized in Table 3 on page 464 of Ravetch and Kinet, *Annu. Rev. Immunol* 9:457-92 (1991).

"Human effector cells" are leukocytes which express one or more FcRs and perform effector functions. Preferably, the cells express at least FcγRIII and perform ADCC effector
30 function. Examples of human leukocytes which mediate ADCC include peripheral blood mononuclear cells (PBMC), natural killer (NK) cells, monocytes, cytotoxic T cells and neutrophils; with PBMCs and NK cells being preferred. The effector cells may be isolated from a native source thereof, *e.g.* from blood or PBMCs as described herein.

A polypeptide variant with "altered" FcR binding affinity or ADCC activity is one which has
35 either enhanced or diminished FcR binding activity and/or ADCC activity compared to a parent

polypeptide or to a polypeptide comprising a native sequence Fc region. The polypeptide variant which "displays increased binding" to an FcR binds at least one FcR with better affinity than the parent polypeptide. The polypeptide variant which "displays decreased binding" to an FcR, binds at least one FcR with worse affinity than a parent polypeptide. Such variants which display
5 decreased binding to an FcR may possess little or no appreciable binding to an FcR, *e.g.*, 0-20% binding to the FcR compared to a native sequence IgG Fc region, *e.g.* as determined in the Examples herein.

The polypeptide variant which binds an FcR with "better affinity" than a parent polypeptide, is one which binds any one or more of the above identified FcRs with substantially better binding
10 affinity than the parent antibody, when the amounts of polypeptide variant and parent polypeptide in the binding assay are essentially the same. For example, the polypeptide variant with improved FcR binding affinity may display from about 1.15 fold to about 100 fold, *e.g.* from about 1.2 fold to about 50 fold improvement in FcR binding affinity compared to the parent polypeptide, where FcR binding affinity is determined, for example, as disclosed in the Examples herein.

The polypeptide variant which "mediates antibody-dependent cell-mediated cytotoxicity (ADCC) in the presence of human effector cells more effectively" than a parent antibody is one which *in vitro* or *in vivo* is substantially more effective at mediating ADCC, when the amounts of polypeptide variant and parent antibody used in the assay are essentially the same. Generally,
15 such variants will be identified using the *in vitro* ADCC assay as herein disclosed, but other assays or methods for determining ADCC activity, *e.g.* in an animal model etc, are contemplated. The preferred variant is from about 1.5 fold to about 100 fold, *e.g.* from about two fold to about fifty fold, more effective at mediating ADCC than the parent, *e.g.* in the *in vitro* assay disclosed herein.

An "amino acid modification" refers to a change in the amino acid sequence of a predetermined amino acid sequence. Exemplary modifications include an amino acid substitution,
25 insertion and/or deletion. The preferred amino acid modification herein is a substitution.

An "amino acid modification at" a specified position, *e.g.* of the Fc region, refers to the substitution or deletion of the specified residue, or the insertion of at least one amino acid residue adjacent the specified residue. By insertion "adjacent" a specified residue is meant insertion
30 within one to two residues thereof. The insertion may be N-terminal or C-terminal to the specified residue.

An "amino acid substitution" refers to the replacement of at least one existing amino acid residue in a predetermined amino acid sequence with another different "replacement" amino acid residue. The replacement residue or residues may be "naturally occurring amino acid residues"
35 (*i.e.* encoded by the genetic code) and selected from the group consisting of: alanine (Ala);

arginine (Arg); asparagine (Asn); aspartic acid (Asp); cysteine (Cys); glutamine (Gln); glutamic acid (Glu); glycine (Gly); histidine (His); isoleucine (Ile); leucine (Leu); lysine (Lys); methionine (Met); phenylalanine (Phe); proline (Pro); serine (Ser); threonine (Thr); tryptophan (Trp); tyrosine (Tyr); and valine (Val). Preferably, the replacement residue is not cysteine. Substitution with one or more non-naturally occurring amino acid residues is also encompassed by the definition of an amino acid substitution herein. A "non-naturally occurring amino acid residue" refers to a residue, other than those naturally occurring amino acid residues listed above, which is able to covalently bind adjacent amino acid residues(s) in a polypeptide chain. Examples of non-naturally occurring amino acid residues include norleucine, ornithine, norvaline, homoserine and other amino acid residue analogues such as those described in Ellman *et al. Meth. Enzym.* 202:301-336 (1991). To generate such non-naturally occurring amino acid residues, the procedures of Noren *et al. Science* 244:182 (1989) and Ellman *et al., supra*, can be used. Briefly, these procedures involve chemically activating a suppressor tRNA with a non-naturally occurring amino acid residue followed by *in vitro* transcription and translation of the RNA.

An "amino acid insertion" refers to the incorporation of at least one amino acid into a predetermined amino acid sequence. While the insertion will usually consist of the insertion of one or two amino acid residues, the present application contemplates larger "peptide insertions", *e.g.* insertion of about three to about five or even up to about ten amino acid residues. The inserted residue(s) may be naturally occurring or non-naturally occurring as disclosed above.

An "amino acid deletion" refers to the removal of at least one amino acid residue from a predetermined amino acid sequence.

"Hinge region" is generally defined as stretching from Glu216 to Pro230 of human IgG1 (Burton, *Molec. Immunol.* 22:161-206 (1985)). Hinge regions of other IgG isotypes may be aligned with the IgG1 sequence by placing the first and last cysteine residues forming inter-heavy chain S-S bonds in the same positions.

The "lower hinge region" of an Fc region is normally defined as the stretch of residues immediately C-terminal to the hinge region, *i.e.* residues 233 to 239 of the Fc region. Prior to the present invention, Fc γ R binding was generally attributed to amino acid residues in the lower hinge region of an IgG Fc region.

"C1q" is a polypeptide that includes a binding site for the Fc region of an immunoglobulin. C1q together with two serine proteases, C1r and C1s, forms the complex C1, the first component of the complement dependent cytotoxicity (CDC) pathway. Human C1q can be purchased commercially from, *e.g.* Quidel, San Diego, CA.

The term "binding domain" refers to the region of a polypeptide that binds to another molecule. In the case of an FcR, the binding domain can comprise a portion of a polypeptide

chain thereof (e.g. the α chain thereof) which is responsible for binding an Fc region. One useful binding domain is the extracellular domain of an FcR α chain.

The term "antibody" is used in the broadest sense and specifically covers monoclonal antibodies (including full length monoclonal antibodies), polyclonal antibodies, multispecific
5 antibodies (e.g., bispecific antibodies), and antibody fragments so long as they exhibit the desired biological activity.

"Antibody fragments", as defined for the purpose of the present invention, comprise a portion of an intact antibody, generally including the antigen binding or variable region of the intact antibody or the Fc region of an antibody which retains FcR binding capability. Examples of
10 antibody fragments include linear antibodies; single-chain antibody molecules; and multispecific antibodies formed from antibody fragments. The antibody fragments preferably retain at least part of the hinge and optionally the CH1 region of an IgG heavy chain. More preferably, the antibody fragments retain the entire constant region of an IgG heavy chain, and include an IgG light chain.

The term "monoclonal antibody" as used herein refers to an antibody obtained from a
15 population of substantially homogeneous antibodies, i.e., the individual antibodies comprising the population are identical except for possible naturally occurring mutations that may be present in minor amounts. Monoclonal antibodies are highly specific, being directed against a single antigenic site. Furthermore, in contrast to conventional (polyclonal) antibody preparations that typically include different antibodies directed against different determinants (epitopes), each
20 monoclonal antibody is directed against a single determinant on the antigen. The modifier "monoclonal" indicates the character of the antibody as being obtained from a substantially homogeneous population of antibodies, and is not to be construed as requiring production of the antibody by any particular method. For example, the monoclonal antibodies to be used in accordance with the present invention may be made by the hybridoma method first described by
25 Kohler *et al.*, *Nature* 256:495 (1975), or may be made by recombinant DNA methods (see, e.g., U.S. Patent No. 4,816,567). The "monoclonal antibodies" may also be isolated from phage antibody libraries using the techniques described in Clackson *et al.*, *Nature* 352:624-628 (1991) and Marks *et al.*, *J. Mol. Biol.* 222:581-597 (1991), for example.

The monoclonal antibodies herein specifically include "chimeric" antibodies
30 (immunoglobulins) in which a portion of the heavy and/or light chain is identical with or homologous to corresponding sequences in antibodies derived from a particular species or belonging to a particular antibody class or subclass, while the remainder of the chain(s) is identical with or homologous to corresponding sequences in antibodies derived from another species or belonging to another antibody class or subclass, as well as fragments of such

antibodies, so long as they exhibit the desired biological activity (U.S. Patent No. 4,816,567; and Morrison *et al.*, *Proc. Natl. Acad. Sci. USA* 81:6851-6855 (1984)).

"Humanized" forms of non-human (*e.g.*, murine) antibodies are chimeric antibodies that contain minimal sequence derived from non-human immunoglobulin. For the most part, humanized antibodies are human immunoglobulins (recipient antibody) in which residues from a hypervariable region of the recipient are replaced by residues from a hypervariable region of a non-human species (donor antibody) such as mouse, rat, rabbit or nonhuman primate having the desired specificity, affinity, and capacity. In some instances, Fv framework region (FR) residues of the human immunoglobulin are replaced by corresponding non-human residues. Furthermore, humanized antibodies may comprise residues that are not found in the recipient antibody or in the donor antibody. These modifications are made to further refine antibody performance. In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the hypervariable loops correspond to those of a non-human immunoglobulin and all or substantially all of the FR regions are those of a human immunoglobulin sequence. The humanized antibody optionally also will comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin. For further details, see Jones *et al.*, *Nature* 321:522-525 (1986); Riechmann *et al.*, *Nature* 332:323-329 (1988); and Presta, *Curr. Op. Struct. Biol.* 2:593-596 (1992).

The term "hypervariable region" when used herein refers to the amino acid residues of an antibody which are responsible for antigen-binding. The hypervariable region comprises amino acid residues from a "complementarity determining region" or "CDR" (*i.e.* residues 24-34 (L1), 50-56 (L2) and 89-97 (L3) in the light chain variable domain and 31-35 (H1), 50-65 (H2) and 95-102 (H3) in the heavy chain variable domain; Kabat *et al.*, *Sequences of Proteins of Immunological Interest*, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, MD. (1991)) and/or those residues from a "hypervariable loop" (*i.e.* residues 26-32 (L1), 50-52 (L2) and 91-96 (L3) in the light chain variable domain and 26-32 (H1), 53-55 (H2) and 96-101 (H3) in the heavy chain variable domain; Chothia and Lesk *J. Mol. Biol.* 196:901-917 (1987)). "Framework" or "FR" residues are those variable domain residues other than the hypervariable region residues as herein defined.

As used herein, the term "immunoadhesin" designates antibody-like molecules which combine the "binding domain" of a heterologous "adhesin" protein (*e.g.* a receptor, ligand or enzyme) with an immunoglobulin constant domain. Structurally, the immunoadhesins comprise a fusion of the adhesin amino acid sequence with the desired binding specificity which is other than the antigen recognition and binding site (antigen combining site) of an antibody (*i.e.* is "heterologous") and an immunoglobulin constant domain sequence.

5 The term "ligand binding domain" as used herein refers to any native cell-surface receptor or any region or derivative thereof retaining at least a qualitative ligand binding ability of a corresponding native receptor. In a specific embodiment, the receptor is from a cell-surface polypeptide having an extracellular domain that is homologous to a member of the immunoglobulin supergenefamily. Other receptors, which are not members of the immunoglobulin supergenefamily but are nonetheless specifically covered by this definition, are receptors for cytokines, and in particular receptors with tyrosine kinase activity (receptor tyrosine kinases), members of the hematopoietin and nerve growth factor receptor superfamilies, and cell adhesion molecules, *e.g.* (E-, L- and P-) selectins.

10 The term "receptor binding domain" is used to designate any native ligand for a receptor, including cell adhesion molecules, or any region or derivative of such native ligand retaining at least a qualitative receptor binding ability of a corresponding native ligand. This definition, among others, specifically includes binding sequences from ligands for the above-mentioned receptors.

15 An "antibody-immunoadhesin chimera" comprises a molecule that combines at least one binding domain of an antibody (as herein defined) with at least one immunoadhesin (as defined in this application). Exemplary antibody-immunoadhesin chimeras are the bispecific CD4-IgG chimeras described in Berg *et al.*, *PNAS (USA)* 88:4723-4727 (1991) and Chamow *et al.*, *J. Immunol.* 153:4268 (1994).

20 An "isolated" polypeptide is one that has been identified and separated and/or recovered from a component of its natural environment. Contaminant components of its natural environment are materials that would interfere with diagnostic or therapeutic uses for the polypeptide, and may include enzymes, hormones, and other proteinaceous or nonproteinaceous solutes. In preferred embodiments, the polypeptide will be purified (1) to greater than 95% by weight of polypeptide as determined by the Lowry method, and most preferably more than 99% by weight, (2) to a degree sufficient to obtain at least 15 residues of N-terminal or internal amino acid sequence by use of a spinning cup sequenator, or (3) to homogeneity by SDS-PAGE under reducing or nonreducing conditions using Coomassie blue or, preferably, silver stain. Isolated polypeptide includes the polypeptide *in situ* within recombinant cells since at least one component of the polypeptide's natural environment will not be present. Ordinarily, however, isolated polypeptide will be prepared by at least one purification step.

30 "Treatment" refers to both therapeutic treatment and prophylactic or preventative measures. Those in need of treatment include those already with the disorder as well as those in which the disorder is to be prevented.

35 A "disorder" is any condition that would benefit from treatment with the polypeptide variant. This includes chronic and acute disorders or diseases including those pathological conditions

which predispose the mammal to the disorder in question. In one embodiment, the disorder is cancer.

The terms "cancer" and "cancerous" refer to or describe the physiological condition in mammals that is typically characterized by unregulated cell growth. Examples of cancer include but are not limited to, carcinoma, lymphoma, blastoma, sarcoma, and leukemia. More particular examples of such cancers include squamous cell cancer, small-cell lung cancer, non-small cell lung cancer, adenocarcinoma of the lung, squamous carcinoma of the lung, cancer of the peritoneum, hepatocellular cancer, gastrointestinal cancer, pancreatic cancer, glioblastoma, cervical cancer, ovarian cancer, liver cancer, bladder cancer, hepatoma, breast cancer, colon cancer, colorectal cancer, endometrial or uterine carcinoma, salivary gland carcinoma, kidney cancer, liver cancer, prostate cancer, vulval cancer, thyroid cancer, hepatic carcinoma and various types of head and neck cancer.

A "HER2-expressing cancer" is one comprising cells which have HER2 receptor protein (Semba *et al.*, *PNAS (USA)* 82:6497-6501 (1985) and Yamamoto *et al.* *Nature* 319:230-234 (1986) (Genebank accession number X03363)) present at their cell surface, such that an anti-HER2 antibody is able to bind to the cancer.

The word "label" when used herein refers to a detectable compound or composition which is conjugated directly or indirectly to the polypeptide. The label may be itself be detectable (e.g., radioisotope labels or fluorescent labels) or, in the case of an enzymatic label, may catalyze chemical alteration of a substrate compound or composition which is detectable.

An "isolated" nucleic acid molecule is a nucleic acid molecule that is identified and separated from at least one contaminant nucleic acid molecule with which it is ordinarily associated in the natural source of the polypeptide nucleic acid. An isolated nucleic acid molecule is other than in the form or setting in which it is found in nature. Isolated nucleic acid molecules therefore are distinguished from the nucleic acid molecule as it exists in natural cells. However, an isolated nucleic acid molecule includes a nucleic acid molecule contained in cells that ordinarily express the polypeptide where, for example, the nucleic acid molecule is in a chromosomal location different from that of natural cells.

The expression "control sequences" refers to DNA sequences necessary for the expression of an operably linked coding sequence in a particular host organism. The control sequences that are suitable for prokaryotes, for example, include a promoter, optionally an operator sequence, and a ribosome binding site. Eukaryotic cells are known to utilize promoters, polyadenylation signals, and enhancers.

Nucleic acid is "operably linked" when it is placed into a functional relationship with another nucleic acid sequence. For example, DNA for a presequence or secretory leader is

operably linked to DNA for a polypeptide if it is expressed as a preprotein that participates in the secretion of the polypeptide; a promoter or enhancer is operably linked to a coding sequence if it affects the transcription of the sequence; or a ribosome binding site is operably linked to a coding sequence if it is positioned so as to facilitate translation. Generally, "operably linked" means that the DNA sequences being linked are contiguous, and, in the case of a secretory leader, contiguous and in reading phase. However, enhancers do not have to be contiguous. Linking is accomplished by ligation at convenient restriction sites. If such sites do not exist, the synthetic oligonucleotide adaptors or linkers are used in accordance with conventional practice.

As used herein, the expressions "cell," "cell line," and "cell culture" are used interchangeably and all such designations include progeny. Thus, the words "transformants" and "transformed cells" include the primary subject cell and cultures derived therefrom without regard for the number of transfers. It is also understood that all progeny may not be precisely identical in DNA content, due to deliberate or inadvertent mutations. Mutant progeny that have the same function or biological activity as screened for in the originally transformed cell are included. Where distinct designations are intended, it will be clear from the context.

The term "molecular complex" when used herein refers to the relatively stable structure which forms when two or more heterologous molecules (*e.g.* polypeptides) bind (preferably noncovalently) to one another. The preferred molecular complex herein is an immune complex.

"Immune complex" refers to the relatively stable structure which forms when at least one target molecule and at least one heterologous Fc region-containing polypeptide bind to one another forming a larger molecular weight complex. Examples of immune complexes are antigen-antibody aggregates and target molecule-immunoadhesin aggregates. The term "immune complex" as used herein, unless indicated otherwise, refers to an *ex vivo* complex (*i.e.* other than the form or setting in which it may be found in nature). However, the immune complex may be administered to a mammal, *e.g.* to evaluate clearance of the immune complex in the mammal.

The term "target molecule" refers to a molecule, usually a polypeptide, which is capable of being bound by a heterologous molecule and has one or more binding sites for the heterologous molecule. The term "binding site" refers to a region of a molecule to which another molecule can bind. The "first target molecule" herein comprises at least two distinct binding sites (for example, two to five separate binding sites) for an analyte (*e.g.* an Fc region-containing polypeptide) such that at least two analyte molecules can bind to the first target molecule. In the preferred embodiment of the invention, the two or more binding sites are identical (*e.g.* having the same amino acid sequence, where the target molecule is a polypeptide). In Example 1 below, the first target molecule was IgE and had two separate binding sites in the Fc region thereof to which the Fc region-containing polypeptide (an anti-IgE antibody, E27) could bind. Other first

target molecules include dimers of substantially identical monomers (*e.g.* neurotrophins, IL8 and VEGF) or are polypeptides comprising two or more substantially identical polypeptide chains (*e.g.* antibodies or immunoadhesins). The "second target molecule" comprises at least two distinct binding sites (for example, two to five separate binding sites) for the first target molecule such that at least two first target molecules can bind to the second target molecule. Preferably, the two or more binding sites are identical (*e.g.* having the same amino acid sequence, where the target molecule is a polypeptide). In Example 2, the second target molecule was VEGF, which has a pair of distinct binding sites to which the variable domain of the IgE antibody could bind. Other second target molecules are contemplated, *e.g.* other dimers of substantially identical monomers (*e.g.* neurotrophins or IL8) or polypeptides comprising two or more substantially identical domains (*e.g.* antibodies or immunoadhesins).

An "analyte" is a substance that is to be analyzed. The preferred analyte is an Fc region-containing polypeptide that is to be analyzed for its ability to bind to an Fc receptor.

A "receptor" is a polypeptide capable of binding at least one ligand. The preferred receptor is a cell-surface receptor having an extracellular ligand-binding domain and, optionally, other domains (*e.g.* transmembrane domain, intracellular domain and/or membrane anchor). The receptor to be evaluated in the assay described herein may be an intact receptor or a fragment or derivative thereof (*e.g.* a fusion protein comprising the binding domain of the receptor fused to one or more heterologous polypeptides). Moreover, the receptor to be evaluated for its binding properties may be present in a cell or isolated and optionally coated on an assay plate or some other solid phase.

The phrase "low affinity receptor" denotes a receptor that has a weak binding affinity for a ligand of interest, *e.g.* having a binding constant of about 50nM or worse affinity. Exemplary low affinity receptors include FcγRII and FcγRIII.

II. Modes for Carrying Out the Invention

The invention herein relates to a method for making a polypeptide variant. The "parent", "starting" or "nonvariant" polypeptide is prepared using techniques available in the art for generating polypeptides comprising an Fc region. In the preferred embodiment of the invention, the parent polypeptide is an antibody and exemplary methods for generating antibodies are described in more detail in the following sections. The parent polypeptide may, however, be any other polypeptide comprising an Fc region, *e.g.* an immunoadhesin. Methods for making immunoadhesins are elaborated in more detail hereinbelow.

In an alternative embodiment, a variant Fc region may be generated according to the methods herein disclosed and this "variant Fc region" can be fused to a heterologous polypeptide of choice, such as an antibody variable domain or binding domain of a receptor or ligand.

The parent polypeptide comprises an Fc region. Generally the Fc region of the parent polypeptide will comprise a native sequence Fc region, and preferably a human native sequence Fc region. However, the Fc region of the parent polypeptide may have one or more pre-existing amino acid sequence alterations or modifications from a native sequence Fc region. For example, the C1q binding activity of the Fc region may have been previously altered (other types of Fc region modifications are described in more detail below). In a further embodiment the parent polypeptide Fc region is "conceptual" and, while it does not physically exist, the antibody engineer may decide upon a desired variant Fc region amino acid sequence and generate a polypeptide comprising that sequence or a DNA encoding the desired variant Fc region amino acid sequence.

In the preferred embodiment of the invention, however, a nucleic acid encoding an Fc region of a parent polypeptide is available and this nucleic acid sequence is altered to generate a variant nucleic acid sequence encoding the Fc region variant.

DNA encoding an amino acid sequence variant of the starting polypeptide is prepared by a variety of methods known in the art. These methods include, but are not limited to, preparation by site-directed (or oligonucleotide-mediated) mutagenesis, PCR mutagenesis, and cassette mutagenesis of an earlier prepared DNA encoding the polypeptide

Site-directed mutagenesis is a preferred method for preparing substitution variants. This technique is well known in the art (see, e.g., Carter *et al.* *Nucleic Acids Res.* 13:4431-4443 (1985) and Kunkel *et al.*, *Proc. Natl. Acad. Sci. USA* 82:488 (1987)). Briefly, in carrying out site-directed mutagenesis of DNA, the starting DNA is altered by first hybridizing an oligonucleotide encoding the desired mutation to a single strand of such starting DNA. After hybridization, a DNA polymerase is used to synthesize an entire second strand, using the hybridized oligonucleotide as a primer, and using the single strand of the starting DNA as a template. Thus, the oligonucleotide encoding the desired mutation is incorporated in the resulting double-stranded DNA.

PCR mutagenesis is also suitable for making amino acid sequence variants of the starting polypeptide. See Higuchi, in *PCR Protocols*, pp.177-183 (Academic Press, 1990); and Vallette *et al.*, *Nuc. Acids Res.* 17:723-733 (1989). Briefly, when small amounts of template DNA are used as starting material in a PCR, primers that differ slightly in sequence from the corresponding region in a template DNA can be used to generate relatively large quantities of a specific DNA fragment that differs from the template sequence only at the positions where the primers differ from the template.

Another method for preparing variants, cassette mutagenesis, is based on the technique described by Wells *et al.*, *Gene* 34:315-323 (1985). The starting material is the plasmid (or other vector) comprising the starting polypeptide DNA to be mutated. The codon(s) in the starting DNA

to be mutated are identified. There must be a unique restriction endonuclease site on each side of the identified mutation site(s). If no such restriction sites exist, they may be generated using the above-described oligonucleotide-mediated mutagenesis method to introduce them at appropriate locations in the starting polypeptide DNA. The plasmid DNA is cut at these sites to linearize it. A double-stranded oligonucleotide encoding the sequence of the DNA between the restriction sites but containing the desired mutation(s) is synthesized using standard procedures, wherein the two strands of the oligonucleotide are synthesized separately and then hybridized together using standard techniques. This double-stranded oligonucleotide is referred to as the cassette. This cassette is designed to have 5' and 3' ends that are compatible with the ends of the linearized plasmid, such that it can be directly ligated to the plasmid. This plasmid now contains the mutated DNA sequence.

Alternatively, or additionally, the desired amino acid sequence encoding a polypeptide variant can be determined, and a nucleic acid sequence encoding such amino acid sequence variant can be generated synthetically.

The amino acid sequence of the parent polypeptide is modified in order to generate a variant Fc region with altered Fc receptor binding affinity or activity *in vitro* and/or *in vivo* and/or altered antibody-dependent cell-mediated cytotoxicity (ADCC) activity *in vitro* and/or *in vivo*.

Generally, the modification entails one or more amino acid substitutions. In one embodiment, the replacement residue does not correspond to a residue present in the same position in any of the native sequence Fc regions in Figure 22A. For example, according to this embodiment of the invention, Pro331 of a human IgG3 or IgG1 Fc region is replaced with a residue other than Ser (the corresponding aligned residue found in native sequence human IgG4).

In one embodiment, the residue in the parent polypeptide which is substituted with a replacement residue is not an alanine and/or is not residue Ala339 of an Fc region. In the case of an amino acid substitution, preferably the residue in the parent polypeptide is replaced with an alanine residue. However, the present invention contemplates replacement of the residue of the parent polypeptide with any other amino acid residue. The substitution may, for example, be a "conservative substitution". Such conservative substitutions are shown in Table 1 under the heading of "preferred substitution". More substantial changes may be achieved by making one or more "exemplary substitutions" which are not the preferred substitution in Table 1.

TABLE 1

Original Residue	Exemplary Substitutions	Preferred Substitution
Ala (A)	val; leu; ile	val
Arg (R)	lys; gln; asn	lys
Asn (N)	gln; his; lys; arg	gln
Asp (D)	glu	glu
Cys (C)	ser	ser
Gln (Q)	asn	asn
Glu (E)	asp	asp
Gly (G)	pro; ala	ala
His (H)	asn; gln; lys; arg	arg
Ile (I)	leu; val; met; ala; phe; norleucine	leu
Leu (L)	norleucine; ile; val; met; ala; phe	ile
Lys (K)	arg; gln; asn	arg
Met (M)	leu; phe; ile	leu
Phe (F)	leu; val; ile; ala; tyr	leu
Pro (P)	ala	ala
Ser (S)	thr	thr
Thr (T)	ser	ser
Trp (W)	tyr; phe	tyr
Tyr (Y)	trp; phe; thr; ser	phe
Val (V)	ile; leu; met; phe; ala; norleucine	leu

Substantial modifications in the biological properties of the Fc region may be accomplished by selecting substitutions that differ significantly in their effect on maintaining (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a sheet or helical conformation, (b) the charge or hydrophobicity of the molecule at the target site, or (c) the bulk of the side chain. Naturally occurring residues are divided into groups based on common side-chain properties:

- (1) hydrophobic: norleucine, met, ala, val, leu, ile;
- (2) neutral hydrophilic: cys, ser, thr;
- (3) acidic: asp, glu;
- (4) basic: asn, gln, his, lys, arg;

(5) residues that influence chain orientation: gly, pro; and

(6) aromatic: trp, tyr, phe.

Non-conservative substitutions will entail exchanging a member of one of these classes for a member of another class. Conservative and non-conservative amino acid substitutions are exemplified in Table 8 hereinbelow.

As is demonstrated in Example 4 herein, one can engineer an Fc region variant with altered binding affinity for one or more FcRs. As was shown in that Example, different classes of Fc region variants can be made *e.g.*, as summarized in the following table. Where the variant Fc region has more than one amino acid substitution, generally, but not necessarily, amino acid substitutions in the same class are combined to achieve the desired result.

TABLE 2
CLASSES OF Fc REGION VARIANTS

Class	FcR binding property	Position of Fc region substitution(s)
1A	reduced binding to all Fc γ R	238, 265, 269, 270, 297*, 327, 329
1B	reduced binding to both Fc γ RII and Fc γ RIII	239, 294, 295, 303, 338, 373, 376, 416, 435
2	improved binding to both Fc γ RII and Fc γ RIII	256, 290, 312, 326, 330, 339#, 378, 430
3	improved binding to Fc γ RII and no effect on Fc γ RIII binding	255, 258, 267, 276, 280, 283, 285, 286, 305, 307, 309, 315, 320, 331, 337, 398
4	improved binding to Fc γ RII and reduced binding to Fc γ RIII	268, 272, 301, 322, 340
5	reduced binding to Fc γ RII and no effect on Fc γ RIII binding	292, 324, 335, 414, 419, 438, 439
6	reduced binding to Fc γ RII and improved binding to Fc γ RIII	298, 333
7	no effect on Fc γ RII binding and reduced binding to Fc γ RIII	248, 249, 252, 254, 278, 289, 293, 296, 338, 382, 388, 389, 434, 437
8	no effect on Fc γ RII binding and improved binding to Fc γ RIII	334, 360

* deglycosylated version

Preferably combined with other Fc region modification(s) (*e.g.* as disclosed herein)

Aside from amino acid substitutions, the present invention contemplates other modifications of the parent Fc region amino acid sequence in order to generate an Fc region variant with altered effector function.

One may, for example, delete one or more amino acid residues of the Fc region in order to reduce binding to an FcR. Generally, one will delete one or more of the Fc region residues identified herein as effecting FcR binding (see Example 4 below) in order to generate such an Fc region variant. Generally, no more than one to about ten Fc region residues will be deleted according to this embodiment of the invention. The Fc region herein comprising one or more amino acid deletions will preferably retain at least about 80%, and preferably at least about 90%, and most preferably at least about 95%, of the parent Fc region or of a native sequence human Fc region.

One may also make amino acid insertion Fc region variants, which variants have altered effector function. For example, one may introduce at least one amino acid residue (e.g. one to two amino acid residues and generally no more than ten residues) adjacent to one or more of the Fc region positions identified herein as impacting FcR binding. By "adjacent" is meant within one to two amino acid residues of a Fc region residue identified herein. Such Fc region variants may display enhanced or diminished FcR binding and/or ADCC activity. In order to generate such insertion variants, one may evaluate a co-crystal structure of a polypeptide comprising a binding region of an FcR (e.g. the extracellular domain of the FcR of interest) and the Fc region into which the amino acid residue(s) are to be inserted (see, for example, Deisenhofer, *Biochemistry* 20(9):2361-2370 (1981); and Burmeister *et al.*, *Nature* 342:379-383, (1994)) in order to rationally design an Fc region variant with, e.g., improved FcR binding ability. Such insertion(s) will generally be made in an Fc region loop, but not in the secondary structure (i.e. in a β -strand) of the Fc region.

By introducing the appropriate amino acid sequence modifications in a parent Fc region, one can generate a variant Fc region which (a) mediates antibody-dependent cell-mediated cytotoxicity (ADCC) in the presence of human effector cells more effectively and/or (b) binds an Fc gamma receptor (Fc γ R) with better affinity than the parent polypeptide. Such Fc region variants will generally comprise at least one amino acid modification in the Fc region. Combining amino acid modifications is thought to be particularly desirable. For example, the variant Fc region may include two, three, four, five, etc substitutions therein, e.g. of the specific Fc region positions identified herein.

Preferably, the parent polypeptide Fc region is a human Fc region, e.g. a native sequence human Fc region human IgG1 (A and non-A allotypes), IgG2, IgG3 or IgG4 Fc region. Such sequences are shown in Fig. 23.

To generate an Fc region with improved ADCC activity, the parent polypeptide preferably has pre-existing ADCC activity, e.g., it comprises a human IgG1 or human IgG3 Fc region. In one embodiment, the variant with improved ADCC mediates ADCC substantially more effectively than

an antibody with a native sequence IgG1 or IgG3 Fc region and the antigen-binding region of the variant. Preferably, the variant comprises, or consists essentially of, substitutions of two or three of the residues at positions 298, 333 and 334 of the Fc region. Most preferably, residues at positions 298, 333 and 334 are substituted, (e.g. with alanine residues). Moreover, in order to generate the Fc region variant with improved ADCC activity, one will generally engineer an Fc region variant with improved binding affinity for Fc γ RIII, which is thought to be an important FcR for mediating ADCC. For example, one may introduce an amino acid modification (e.g. a substitution) into the parent Fc region at any one or more of amino acid positions 256, 290, 298, 312, 326, 330, 333, 334, 360, 378 or 430 to generate such a variant. The variant with improved binding affinity for Fc γ RIII may further have reduced binding affinity for Fc γ RII, especially reduced affinity for the inhibiting Fc γ RIIB receptor.

The amino acid modification(s) are preferably introduced into the CH2 domain of a Fc region, since the experiments herein indicate that the CH2 domain is important for FcR binding activity. Moreover, unlike the teachings of the above-cited art, the instant application contemplates the introduction of a modification into a part of the Fc region other than in the lower hinge region thereof.

Useful amino acid positions for modification in order to generate a variant IgG Fc region with altered Fc gamma receptor (Fc γ R) binding affinity or activity include any one or more of amino acid positions 238, 239, 248, 249, 252, 254, 255, 256, 258, 265, 267, 268, 269, 270, 272, 276, 278, 280, 283, 285, 286, 289, 290, 292, 293, 294, 295, 296, 298, 301, 303, 305, 307, 309, 312, 315, 320, 322, 324, 326, 327, 329, 330, 331, 333, 334, 335, 337, 338, 340, 360, 373, 376, 378, 382, 388, 389, 398, 414, 416, 419, 430, 434, 435, 437, 438 or 439 of the Fc region. Preferably, the parent Fc region used as the template to generate such variants comprises a human IgG Fc region. Where residue 331 is substituted, the parent Fc region is preferably not human native sequence IgG3, or the variant Fc region comprising a substitution at position 331 preferably displays increased FcR binding, e.g. to Fc γ RII.

To generate an Fc region variant with reduced binding to the Fc γ R one may introduce an amino acid modification at any one or more of amino acid positions 238, 239, 248, 249, 252, 254, 265, 268, 269, 270, 272, 278, 289, 292, 293, 294, 295, 296, 298, 301, 303, 322, 324, 327, 329, 333, 335, 338, 340, 373, 376, 382, 388, 389, 414, 416, 419, 434, 435, 437, 438 or 439 of the Fc region.

Variants which display reduced binding to Fc γ RI, include those comprising an Fc region amino acid modification at any one or more of amino acid positions 238, 265, 269, 270, 327 or 329.

Variants which display reduced binding to FcγRII include those comprising an Fc region amino acid modification at any one or more of amino acid positions 238, 265, 269, 270, 292, 294, 295, 298, 303, 324, 327, 329, 333, 335, 338, 373, 376, 414, 416, 419, 435, 438 or 439.

5 Fc region variants which display reduced binding to FcγRIII include those comprising an Fc region amino acid modification at any one or more of amino acid positions 238, 239, 248, 249, 252, 254, 265, 268, 269, 270, 272, 278, 289, 293, 294, 295, 296, 301, 303, 322, 327, 329, 338, 340, 373, 376, 382, 388, 389, 416, 434, 435 or 437 .

Variants with improved binding to one or more FcγRs may also be made. Such Fc region variants may comprise an amino acid modification at any one or more of amino acid positions
10 255, 256, 258, 267, 268, 272, 276, 280, 283, 285, 286, 290, 298, 301, 305, 307, 309, 312, 315, 320, 322, 326, 330, 331, 333, 334, 337, 340, 360, 378, 398 or 430 of the Fc region.

For example, the variant with improved FcγR binding activity may display increased binding to FcγRIII, and optionally may further display decreased binding to FcγRII; *e.g.* the variant may comprise an amino acid modification at position 298 and/or 333 of an Fc region.

15 Variants with increased binding to FcγRII include those comprising an amino acid modification at any one or more of amino acid positions 255, 256, 258, 267, 268, 272, 276, 280, 283, 285, 286, 290, 301, 305, 307, 309, 312, 315, 320, 322, 326, 330, 331, 337, 340, 378, 398 or 430 of an Fc region. Such variants may further display decreased binding to FcγRIII. For example, they may include an Fc region amino acid modification at any one or more of amino acid
20 positions 268, 272, 298, 301, 322 or 340.

While it is preferred to alter binding to a FcγR, Fc region variants with altered binding affinity for the neonatal receptor (FcRn) are also contemplated herein. Fc region variants with improved affinity for FcRn are anticipated to have longer serum half-lives, and such molecules will have useful applications in methods of treating mammals where long half-life of the administered polypeptide is desired, *e.g.*, to treat a chronic disease or disorder. Fc region variants with
25 decreased FcRn binding affinity, on the contrary, are expected to have shorter half-lives, and such molecules may, for example, be administered to a mammal where a shortened circulation time may be advantageous, *e.g.* for *in vivo* diagnostic imaging or for polypeptides which have toxic side effects when left circulating in the blood stream for extended periods, etc. Fc region variants
30 with decreased FcRn binding affinity are anticipated to be less likely to cross the placenta, and thus may be utilized in the treatment of diseases or disorders in pregnant women.

Fc region variants with altered binding affinity for FcRn include those comprising an Fc region amino acid modification at any one or more of amino acid positions 238, 252, 253, 254, 255, 256, 265, 272, 286, 288, 303, 305, 307, 309, 311, 312, 317, 340, 356, 360, 362, 376, 378,
35 380, 382, 386, 388, 400, 413, 415, 424, 433, 434, 435, 436, 439 or 447. Those which display

reduced binding to FcRn will generally comprise an Fc region amino acid modification at any one or more of amino acid positions 252, 253, 254, 255, 288, 309, 386, 388, 400, 415, 433, 435, 436, 439 or 447; and those with increased binding to FcRn will usually comprise an Fc region amino acid modification at any one or more of amino acid positions 238, 256, 265, 272, 286, 303, 305, 307, 311, 312, 317, 340, 356, 360, 362, 376, 378, 380, 382, 413, 424 or 434.

The polypeptide variant(s) prepared as described above may be subjected to further modifications, oftentimes depending on the intended use of the polypeptide. Such modifications may involve further alteration of the amino acid sequence (substitution, insertion and/or deletion of amino acid residues), fusion to heterologous polypeptide(s) and/or covalent modifications. Such "further modifications" may be made prior to, simultaneously with, or following, the amino acid modification(s) disclosed above which result in an alteration of Fc receptor binding and/or ADCC activity. In one embodiment, one may combine the Fc region modification herein with Fc region substitutions disclosed in the references cited in the "Related Art" section of this application.

Alternatively or additionally, it may be useful to combine the above amino acid modifications with one or more further amino acid modifications that alter C1q binding and/or complement dependent cytotoxicity function of the Fc region.

The starting polypeptide of particular interest herein is usually one that binds to C1q and displays complement dependent cytotoxicity (CDC). The further amino acid substitutions described herein will generally serve to alter the ability of the starting polypeptide to bind to C1q and/or modify its complement dependent cytotoxicity function, e.g. to reduce and preferably abolish these effector functions. However, polypeptides comprising substitutions at one or more of the described positions with improved C1q binding and/or complement dependent cytotoxicity (CDC) function are contemplated herein. For example, the starting polypeptide may be unable to bind C1q and/or mediate CDC and may be modified according to the teachings herein such that it acquires these further effector functions. Moreover, polypeptides with pre-existing C1q binding activity, optionally further having the ability to mediate CDC may be modified such that one or both of these activities are enhanced.

To generate an Fc region with altered C1q binding and/or complement dependent cytotoxicity (CDC) function, the amino acid positions to be modified are generally selected from heavy chain positions 270, 322, 326, 327, 329, 331, 333, and 334, where the numbering of the residues in an IgG heavy chain is that of the EU index as in Kabat *et al.*, ***Sequences of Proteins of Immunological Interest***, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, MD (1991). In one embodiment, only one of the eight above-identified positions is altered in order to generate the polypeptide variant region with altered C1q binding and/or

complement dependent cytotoxicity (CDC) function. Preferably only residue 270, 329 or 322 is altered if this is the case. Alternatively, two or more of the above-identified positions are modified.

If substitutions are to be combined, generally substitutions which enhance human C1q binding (e.g. at residue positions 326, 327, 333 and 334) or those which diminish human C1q binding (e.g., at residue positions 270, 322, 329 and 331) are combined. In the latter embodiment, all four positions (i.e., 270, 322, 329 and 331) may be substituted. Preferably, further substitutions at two, three or all of positions 326, 327, 333 or 334 are combined, optionally with other Fc region substitutions, to generate a polypeptide with improved human C1q binding and preferably improved CDC activity *in vitro* or *in vivo*.

10 Proline is conserved at position 329 in human IgG's. This residue is preferably replaced with alanine, however substitution with any other amino acid is contemplated, e.g., serine, threonine, asparagine, glycine or valine.

Proline is conserved at position 331 in human IgG1, IgG2 and IgG3, but not IgG4 (which has a serine residue at position 331). Residue 331 is preferably replaced by alanine or another amino acid, e.g. serine (for IgG regions other than IgG4), glycine or valine.

15 Lysine 322 is conserved in human IgGs, and this residue is preferably replaced by an alanine residue, but substitution with any other amino acid residue is contemplated, e.g. serine, threonine, glycine or valine.

D270 is conserved in human IgGs, and this residue may be replaced by another amino acid residue, e.g. alanine, serine, threonine, glycine, valine, or lysine.

K326 is also conserved in human IgGs. This residue may be substituted with another residue including, but not limited to, valine, glutamic acid, alanine, glycine, aspartic acid, methionine or tryptophan, with tryptophan being preferred.

Likewise, E333 is also conserved in human IgGs. E333 is preferably replaced by an amino acid residue with a smaller side chain volume, such as valine, glycine, alanine or serine, with serine being preferred.

K334 is conserved in human IgGs and may be substituted with another residue such as alanine or other residue.

In human IgG1 and IgG3, residue 327 is an alanine. In order to generate a variant with improved C1q binding, this alanine may be substituted with another residue such as glycine. In IgG2 and IgG4, residue 327 is a glycine and this may be replaced by alanine (or another residue) to diminish C1q binding.

As disclosed above, one can design an Fc region with altered effector function, e.g., by modifying C1q binding and/or FcR binding and thereby changing CDC activity and/or ADCC activity. For example, one can generate a variant Fc region with improved C1q binding and

improved FcγRIII binding; *e.g.* having both improved ADCC activity and improved CDC activity. Alternatively, where one desires that effector function be reduced or ablated, one may engineer a variant Fc region with reduced CDC activity and/or reduced ADCC activity. In other embodiments, one may increase only one of these activities, and optionally also reduce the other activity, *e.g.* to generate an Fc region variant with improved ADCC activity, but reduced CDC activity *and vice versa*.

With respect to further amino acid sequence alterations, any cysteine residue not involved in maintaining the proper conformation of the polypeptide variant also may be substituted, generally with serine, to improve the oxidative stability of the molecule and prevent aberrant cross linking.

Another type of amino acid substitution serves to alter the glycosylation pattern of the polypeptide. This may be achieved by deleting one or more carbohydrate moieties found in the polypeptide, and/or adding one or more glycosylation sites that are not present in the polypeptide.

Glycosylation of polypeptides is typically either N-linked or O-linked. N-linked refers to the attachment of the carbohydrate moiety to the side chain of an asparagine residue. The tripeptide sequences asparagine-X-serine and asparagine-X-threonine, where X is any amino acid except proline, are the recognition sequences for enzymatic attachment of the carbohydrate moiety to the asparagine side chain. Thus, the presence of either of these tripeptide sequences in a polypeptide creates a potential glycosylation site. O-linked glycosylation refers to the attachment of one of the sugars N-aceylgalactosamine, galactose, or xylose to a hydroxyamino acid, most commonly serine or threonine, although 5-hydroxyproline or 5-hydroxylysine may also be used.

Addition of glycosylation sites to the polypeptide is conveniently accomplished by altering the amino acid sequence such that it contains one or more of the above-described tripeptide sequences (for N-linked glycosylation sites). The alteration may also be made by the addition of, or substitution by, one or more serine or threonine residues to the sequence of the original polypeptide (for O-linked glycosylation sites). An exemplary glycosylation variant has an amino acid substitution of residue Asn 297 of the heavy chain.

Moreover, the class, subclass or allotype of the Fc region may be altered by one or more further amino acid substitutions to generate an Fc region with an amino acid sequence more homologous to a different class, subclass or allotype as desired. For example, a murine Fc region may be altered to generate an amino acid sequence more homologous to a human Fc region; a human non-A allotype IgG1 Fc region may be modified to achieve a human A allotype IgG1 Fc region etc. In one embodiment, the amino modification(s) herein which alter FcR binding and/or ADCC activity are made in the CH2 domain of the Fc region and the CH3 domain is deleted or

replaced with another dimerization domain. Preferably, however, the CH3 domain is retained (aside from amino acid modifications therein which alter effector function as herein disclosed).

The polypeptide variant may be subjected to one or more assays to evaluate any change in biological activity compared to the starting polypeptide.

5 Preferably the polypeptide variant essentially retains the ability to bind antigen compared to the nonvariant polypeptide, *i.e.* the binding capability is no worse than about 20 fold, *e.g.* no worse than about 5 fold of that of the nonvariant polypeptide. The binding capability of the polypeptide variant may be determined using techniques such as fluorescence activated cell sorting (FACS) analysis or radioimmunoprecipitation (RIA), for example.

10 The ability of the polypeptide variant to bind an FcR may be evaluated. Where the FcR is a high affinity Fc receptor, such as FcγRI, FcRn or FcγRIIIA-V158, binding can be measured by titrating monomeric polypeptide variant and measuring bound polypeptide variant using an antibody which specifically binds to the polypeptide variant in a standard ELISA format (see Example 2 below). Another FcR binding assay for low affinity FcRs is described in Examples 1
15 and 4.

 To assess ADCC activity of the polypeptide variant, an *in vitro* ADCC assay, such as that described in Example 4 may be performed using varying effector:target ratios. Useful "effector cells" for such assays include peripheral blood mononuclear cells (PBMC) and Natural Killer (NK) cells. Alternatively, or additionally, ADCC activity of the polypeptide variant may be *assessed in*
20 *vivo*, *e.g.*, in a animal model such as that disclosed in Clynes *et al. PNAS (USA)* 95:652-656 (1998).

 The ability of the variant to bind C1q and mediate complement dependent cytotoxicity (CDC) may be assessed.

 To determine C1q binding, a C1q binding ELISA may be performed. Briefly, assay plates
25 may be coated overnight at 4°C with polypeptide variant or starting polypeptide (control) in coating buffer. The plates may then be washed and blocked. Following washing, an aliquot of human C1q may be added to each well and incubated for 2 hrs at room temperature. Following a further wash, 100 μl of a sheep anti-complement C1q peroxidase conjugated antibody may be added to each well and incubated for 1 hour at room temperature. The plate may again be washed with
30 wash buffer and 100 μl of substrate buffer containing OPD (O-phenylenediamine dihydrochloride (Sigma)) may be added to each well. The oxidation reaction, observed by the appearance of a yellow color, may be allowed to proceed for 30 minutes and stopped by the addition of 100 μl of 4.5 N H₂SO₄. The absorbance may then read at (492-405) nm.

 An exemplary polypeptide variant is one that displays a "significant reduction in C1q
35 binding" in this assay. This means that about 100μg/ml of the polypeptide variant displays about

50 fold or more reduction in C1q binding compared to 100µg/ml of a control antibody having a nonmutated IgG1 Fc region. In the most preferred embodiment, the polypeptide variant "does not bind C1q", *i.e.* 100µg/ml of the polypeptide variant displays about 100 fold or more reduction in C1q binding compared to 100µg/ml of the control antibody.

5 Another exemplary variant is one which "has a better binding affinity for human C1q than the parent polypeptide". Such a molecule may display, for example, about two-fold or more, and preferably about five-fold or more, improvement in human C1q binding compared to the parent polypeptide (*e.g.* at the IC_{50} values for these two molecules). For example, human C1q binding may be about two-fold to about 500-fold, and preferably from about two-fold or from about five-fold
10 to about 1000-fold improved compared to the parent polypeptide.

To assess complement activation, a complement dependent cytotoxicity (CDC) assay may be performed, *e.g.* as described in Gazzano-Santoro *et al.*, *J. Immunol. Methods* 202:163 (1996). Briefly, various concentrations of the polypeptide variant and human complement may be diluted with buffer. Cells which express the antigen to which the polypeptide variant binds
15 may be diluted to a density of $\sim 1 \times 10^6$ cells /ml. Mixtures of polypeptide variant, diluted human complement and cells expressing the antigen may be added to a flat bottom tissue culture 96 well plate and allowed to incubate for 2 hrs at 37°C and 5% CO₂ to facilitate complement mediated cell lysis. 50 µl of alamar blue (Accumed International) may then be added to each well and incubated overnight at 37°C. The absorbance is measured using a 96-well fluorometer with excitation at
20 530 nm and emission at 590 nm. The results may be expressed in relative fluorescence units (RFU). The sample concentrations may be computed from a standard curve and the percent activity as compared to nonvariant polypeptide is reported for the polypeptide variant of interest.

Yet another exemplary variant "does not activate complement". For example, 0.6 µg/ml of the polypeptide variant displays about 0-10% CDC activity in this assay compared to a 0.6
25 µg/ml of a control antibody having a nonmutated IgG1 Fc region. Preferably the variant does not appear to have any CDC activity in the above CDC assay.

The invention also pertains to a polypeptide variant with enhanced CDC compared to a parent polypeptide, *e.g.*, displaying about two-fold to about 100-fold improvement in CDC activity *in vitro* or *in vivo* (*e.g.* at the IC_{50} values for each molecule being compared).

30 A. Receptor Binding Assay and Immune Complex

A receptor binding assay has been developed herein which is particularly useful for determining binding of an analyte of interest to a receptor where the affinity of the analyte for the receptor is relatively weak, *e.g.* in the micromolar range as is the case for FcγRIIA, FcγRIIB, FcγRIIIA and FcγRIIIB. The method involves the formation of a molecular complex that has an
35 improved avidity for the receptor of interest compared to the noncomplexed analyte. The

preferred molecular complex is an immune complex comprising: (a) an Fc region-containing polypeptide (such as an antibody or an immunoadhesin); (b) a first target molecule which comprises at least two binding sites for the Fc region-containing polypeptide; and (c) a second target molecule which comprises at least two binding sites for the first target molecule.

5 In Example 1 below, the Fc region-containing polypeptide is an anti-IgE antibody, such as the E27 antibody (Figs. 4A-4B). E27, when mixed with human IgE at an 1:1 molar ratio, forms a stable hexamer consisting of three E27 molecules and three IgE molecules. In Example 1 below, the "first target molecule" is a chimeric form of IgE in which the Fab portion of an anti-VEGF antibody is fused to the human IgE Fc portion and the "second target molecule" is the
10 antigen to which the Fab binds (*i.e.* VEGF). Each molecule of IgE binds two molecules of VEGF. VEGF also binds two molecules of IgE per molecule of VEGF. When recombinant human VEGF was added at a 2:1 molar ratio to IgE:E27 hexamers, the hexamers were linked into larger molecular weight complexes via the IgE:VEGF interaction (Fig. 5). The Fc region of the anti-IgE antibody of the resultant immune complex binds to FcR with higher avidity than either
15 uncomplexed anti-IgE or anti-IgE:IgE hexamers.

Other forms of molecular complexes for use in the receptor assay are contemplated. Examples comprising only an Fc region-containing polypeptide:first target molecule combination include an immunoadhesin:ligand combination such as VEGF receptor (KDR)-immunoadhesin:VEGF and a full-length bispecific antibody (bsAb):first target molecule. A further
20 example of an Fc region-containing polypeptide:first target molecule:second target molecule combination include a nonblocking antibody:soluble receptor:ligand combination such as anti-Trk antibody:soluble Trk receptor:neurotrophin (Urfer *et al. J. Biol. Chem.* 273(10):5829-5840 (1998)).

Aside from use in a receptor binding assay, the immune complexes described above
25 have further uses including evaluation of Fc region-containing polypeptide function and immune complex clearance *in vivo*. Hence, the immune complex may be administered to a mammal (*e.g.* in a pre-clinical animal study) and evaluated for its half-life etc.

To determine receptor binding, a polypeptide comprising at least the binding domain of the receptor of interest (*e.g.* the extracellular domain of an α subunit of an FcR) may be coated
30 on solid phase, such as an assay plate. The binding domain of the receptor alone or a receptor-fusion protein may be coated on the plate using standard procedures. Examples of receptor-fusion proteins include receptor-glutathione S-transferase (GST) fusion protein, receptor-chitin binding domain fusion protein, receptor-hexaHis tag fusion protein (coated on glutathione, chitin, and nickel coated plates, respectively). Alternatively, a capture molecule may be coated on the
35 assay plate and used to bind the receptor-fusion protein via the non-receptor portion of the fusion

protein. Examples include anti-hexaHis F(ab')₂ coated on the assay plate used to capture receptor-hexaHis tail fusion or anti-GST antibody coated on the assay plate used to capture a receptor-GST fusion. In other embodiments, binding to cells expressing at least the binding domain of the receptor may be evaluated. The cells may be naturally occurring hematopoietic cells that express the FcR of interest or may be transformed with nucleic acid encoding the FcR or a binding domain thereof such that the binding domain is expressed at the surface of the cell to be tested.

The immune complex described hereinabove is added to the receptor-coated plates and incubated for a sufficient period of time such that the analyte binds to the receptor. Plates may then be washed to remove unbound complexes, and binding of the analyte may be detected according to known methods. For example, binding may be detected using a reagent (e.g. an antibody or fragment thereof) which binds specifically to the analyte, and which is optionally conjugated with a detectable label (detectable labels and methods for conjugating them to polypeptides are described below in the section entitled "Non-Therapeutic Uses for the Polypeptide Variant").

As a matter of convenience, the reagents can be provided in an assay kit, i.e., a packaged combination of reagents, for combination with the analyte in assaying the ability of the analyte to bind to a receptor of interest. The components of the kit will generally be provided in predetermined ratios. The kit may provide the first target molecule and/or the second target molecule, optionally complexed together. The kit may further include assay plates coated with the receptor or a binding domain thereof (e.g. the extracellular domain of the α subunit of an FcR). Usually, other reagents, such as an antibody that binds specifically to the analyte to be assayed, labeled directly or indirectly with an enzymatic label, will also be provided in the kit. Where the detectable label is an enzyme, the kit will include substrates and cofactors required by the enzyme (e.g. a substrate precursor which provides the detectable chromophore or fluorophore). In addition, other additives may be included such as stabilizers, buffers (e.g. assay and/or wash lysis buffer) and the like. The relative amounts of the various reagents may be varied widely to provide for concentrations in solution of the reagents that substantially optimize the sensitivity of the assay. Particularly, the reagents may be provided as dry powders, usually lyophilized, including excipients that on dissolution will provide a reagent solution having the appropriate concentration. The kit also suitably includes instructions for carrying out the assay.

B. Antibody Preparation

In the preferred embodiment of the invention, the Fc region-containing polypeptide which is modified according to the teachings herein is an antibody. Techniques for producing antibodies follow:

(i) Antigen selection and preparation

Where the polypeptide is an antibody, it is directed against an antigen of interest. Preferably, the antigen is a biologically important polypeptide and administration of the antibody to a mammal suffering from a disease or disorder can result in a therapeutic benefit in that mammal. However, antibodies directed against nonpolypeptide antigens (such as tumor-associated glycolipid antigens; see US Patent 5,091,178) are also contemplated.

Where the antigen is a polypeptide, it may be a transmembrane molecule (*e.g.* receptor) or ligand such as a growth factor. Exemplary antigens include molecules such as renin; a growth hormone, including human growth hormone and bovine growth hormone; growth hormone releasing factor; parathyroid hormone; thyroid stimulating hormone; lipoproteins; alpha-1-antitrypsin; insulin A-chain; insulin B-chain; proinsulin; follicle stimulating hormone; calcitonin; luteinizing hormone; glucagon; clotting factors such as factor VIIIc, factor IX, tissue factor (TF), and von Willebrands factor; anti-clotting factors such as Protein C; atrial natriuretic factor; lung surfactant; a plasminogen activator, such as urokinase or human urine or tissue-type plasminogen activator (t-PA); bombesin; thrombin; hemopoietic growth factor; tumor necrosis factor-alpha and -beta; enkephalinase; RANTES (regulated on activation normally T-cell expressed and secreted); human macrophage inflammatory protein (MIP-1-alpha); a serum albumin such as human serum albumin; Muellerian-inhibiting substance; relaxin A-chain; relaxin B-chain; prorelaxin; mouse gonadotropin-associated peptide; a microbial protein, such as beta-lactamase; DNase; IgE; a cytotoxic T-lymphocyte associated antigen (CTLA), such as CTLA-4; inhibin; activin; vascular endothelial growth factor (VEGF); receptors for hormones or growth factors; protein A or D; rheumatoid factors; a neurotrophic factor such as bone-derived neurotrophic factor (BDNF), neurotrophin-3, -4, -5, or -6 (NT-3, NT-4, NT-5, or NT-6), or a nerve growth factor such as NGF- β ; platelet-derived growth factor (PDGF); fibroblast growth factor such as aFGF and bFGF; epidermal growth factor (EGF); transforming growth factor (TGF) such as TGF-alpha and TGF-beta, including TGF- β 1, TGF- β 2, TGF- β 3, TGF- β 4, or TGF- β 5; insulin-like growth factor-I and -II (IGF-I and IGF-II); des(1-3)-IGF-I (brain IGF-I), insulin-like growth factor binding proteins; CD proteins such as CD3, CD4, CD8, CD19 and CD20; erythropoietin; osteoinductive factors; immunotoxins; a bone morphogenetic protein (BMP); an interferon such as interferon-alpha, -beta, and -gamma; colony stimulating factors (CSFs), *e.g.*, M-CSF, GM-CSF, and G-CSF; interleukins (ILs), *e.g.*, IL-1 to IL-10; superoxide dismutase; T-cell receptors; surface membrane proteins; decay accelerating factor; viral antigen such as, for example, a portion of the AIDS envelope; transport proteins; homing receptors; addressins; regulatory proteins; integrins such as CD11a, CD11b, CD11c, CD18, an ICAM, VLA-4 and VCAM; a tumor associated antigen such as HER2, HER3 or HER4 receptor; and fragments of any of the above-listed polypeptides.

Preferred molecular targets for antibodies encompassed by the present invention include CD proteins such as CD3, CD4, CD8, CD19, CD20 and CD34; members of the ErbB receptor family such as the EGF receptor, HER2, HER3 or HER4 receptor; cell adhesion molecules such as LFA-1, Mac1, p150.95, VLA-4, ICAM-1, VCAM, $\alpha 4/\beta 7$ integrin, and $\alpha v/\beta 3$ integrin including
5 either α or β subunits thereof (e.g. anti-CD11a, anti-CD18 or anti-CD11b antibodies); growth factors such as VEGF; tissue factor (TF); alpha interferon (α -IFN); an interleukin, such as IL-8; IgE; blood group antigens; flk2/flt3 receptor; obesity (OB) receptor; *mpl* receptor; CTLA-4; protein C etc.

Soluble antigens or fragments thereof, optionally conjugated to other molecules, can be
10 used as immunogens for generating antibodies. For transmembrane molecules, such as receptors, fragments of these (e.g. the extracellular domain of a receptor) can be used as the immunogen. Alternatively, cells expressing the transmembrane molecule can be used as the immunogen. Such cells can be derived from a natural source (e.g. cancer cell lines) or may be cells which have been transformed by recombinant techniques to express the transmembrane
15 molecule. Other antigens and forms thereof useful for preparing antibodies will be apparent to those in the art.

(ii) *Polyclonal antibodies*

Polyclonal antibodies are preferably raised in animals by multiple subcutaneous (sc) or intraperitoneal (ip) injections of the relevant antigen and an adjuvant. It may be useful to
20 conjugate the relevant antigen to a protein that is immunogenic in the species to be immunized, e.g., keyhole limpet hemocyanin, serum albumin, bovine thyroglobulin, or soybean trypsin inhibitor using a bifunctional or derivatizing agent, for example, maleimidobenzoyl sulfosuccinimide ester (conjugation through cysteine residues), N-hydroxysuccinimide (through lysine residues), glutaraldehyde, succinic anhydride, SOCl_2 , or $\text{R}^1\text{N}=\text{C}=\text{NR}$, where R and R^1 are different alkyl
25 groups.

Animals are immunized against the antigen, immunogenic conjugates, or derivatives by combining, e.g., 100 μg or 5 μg of the protein or conjugate (for rabbits or mice, respectively) with 3 volumes of Freund's complete adjuvant and injecting the solution intradermally at multiple sites. One month later the animals are boosted with 1/5 to 1/10 the original amount of peptide or
30 conjugate in Freund's complete adjuvant by subcutaneous injection at multiple sites. Seven to 14 days later the animals are bled and the serum is assayed for antibody titer. Animals are boosted until the titer plateaus. Preferably, the animal is boosted with the conjugate of the same antigen, but conjugated to a different protein and/or through a different cross-linking reagent. Conjugates also can be made in recombinant cell culture as protein fusions. Also, aggregating
35 agents such as alum are suitably used to enhance the immune response.

(iii) **Monoclonal antibodies**

Monoclonal antibodies may be made using the hybridoma method first described by Kohler *et al.*, *Nature*, 256:495 (1975), or may be made by recombinant DNA methods (U.S. Patent No. 4,816,567).

5 In the hybridoma method, a mouse or other appropriate host animal, such as a hamster or macaque monkey, is immunized as hereinabove described to elicit lymphocytes that produce or are capable of producing antibodies that will specifically bind to the protein used for immunization. Alternatively, lymphocytes may be immunized *in vitro*. Lymphocytes then are fused with myeloma cells using a suitable fusing agent, such as polyethylene glycol, to form a
10 hybridoma cell (Goding, *Monoclonal Antibodies: Principles and Practice*, pp.59-103 (Academic Press, 1986)).

The hybridoma cells thus prepared are seeded and grown in a suitable culture medium that preferably contains one or more substances that inhibit the growth or survival of the unfused, parental myeloma cells. For example, if the parental myeloma cells lack the enzyme
15 hypoxanthine guanine phosphoribosyl transferase (HGPRT or HPRT), the culture medium for the hybridomas typically will include hypoxanthine, aminopterin, and thymidine (HAT medium), which substances prevent the growth of HGPRT-deficient cells.

Preferred myeloma cells are those that fuse efficiently, support stable high-level production of antibody by the selected antibody-producing cells, and are sensitive to a medium
20 such as HAT medium. Among these, preferred myeloma cell lines are murine myeloma lines, such as those derived from MOPC-21 and MPC-11 mouse tumors available from the Salk Institute Cell Distribution Center, San Diego, California USA, and SP-2 or X63-Ag8-653 cells available from the American Type Culture Collection, Rockville, Maryland USA. Human myeloma and mouse-human heteromyeloma cell lines also have been described for the production of human
25 monoclonal antibodies (Kozbor, *J. Immunol.*, 133:3001 (1984); Brodeur *et al.*, *Monoclonal Antibody Production Techniques and Applications*, pp. 51-63 (Marcel Dekker, Inc., New York, 1987)).

Culture medium in which hybridoma cells are growing is assayed for production of monoclonal antibodies directed against the antigen. Preferably, the binding specificity of
30 monoclonal antibodies produced by hybridoma cells is determined by immunoprecipitation or by an *in vitro* binding assay, such as radioimmunoassay (RIA) or enzyme-linked immunoabsorbent assay (ELISA).

After hybridoma cells are identified that produce antibodies of the desired specificity, affinity, and/or activity, the clones may be subcloned by limiting dilution procedures and grown by
35 standard methods (Goding, *Monoclonal Antibodies: Principles and Practice*, pp.59-103

(Academic Press, 1986)). Suitable culture media for this purpose include, for example, D-MEM or RPMI-1640 medium. In addition, the hybridoma cells may be grown *in vivo* as ascites tumors in an animal.

5 The monoclonal antibodies secreted by the subclones are suitably separated from the culture medium, ascites fluid, or serum by conventional immunoglobulin purification procedures such as, for example, protein A-Sepharose, hydroxylapatite chromatography, gel electrophoresis, dialysis, or affinity chromatography.

DNA encoding the monoclonal antibodies is readily isolated and sequenced using conventional procedures (*e.g.*, by using oligonucleotide probes that are capable of binding specifically to genes encoding the heavy and light chains of the monoclonal antibodies). The hybridoma cells serve as a preferred source of such DNA. Once isolated, the DNA may be placed into expression vectors, which are then transfected into host cells such as *E. coli* cells, simian COS cells, Chinese hamster ovary (CHO) cells, or myeloma cells that do not otherwise produce immunoglobulin protein, to obtain the synthesis of monoclonal antibodies in the recombinant host cells. Recombinant production of antibodies will be described in more detail below.

10 In a further embodiment, antibodies or antibody fragments can be isolated from antibody phage libraries generated using the techniques described in McCafferty *et al.*, *Nature*, 348:552-554 (1990). Clackson *et al.*, *Nature*, 352:624-628 (1991) and Marks *et al.*, *J. Mol. Biol.*, 222:581-597 (1991) describe the isolation of murine and human antibodies, respectively, using phage libraries. Subsequent publications describe the production of high affinity (nM range) human antibodies by chain shuffling (Marks *et al.*, *Bio/Technology*, 10:779-783 (1992)), as well as combinatorial infection and *in vivo* recombination as a strategy for constructing very large phage libraries (Waterhouse *et al.*, *Nuc. Acids. Res.*, 21:2265-2266 (1993)). Thus, these techniques are viable alternatives to traditional monoclonal antibody hybridoma techniques for isolation of monoclonal antibodies.

25 The DNA also may be modified, for example, by substituting the coding sequence for human heavy- and light-chain constant domains in place of the homologous murine sequences (U.S. Patent No. 4,816,567; Morrison, *et al.*, *Proc. Natl Acad. Sci. USA*, 81:6851 (1984)), or by covalently joining to the immunoglobulin coding sequence all or part of the coding sequence for a non-immunoglobulin polypeptide.

30 Typically such non-immunoglobulin polypeptides are substituted for the constant domains of an antibody, or they are substituted for the variable domains of one antigen-combining site of an antibody to create a chimeric bivalent antibody comprising one antigen-combining site having specificity for an antigen and another antigen-combining site having specificity for a different antigen.

(iv) **Humanized and human antibodies**

A humanized antibody has one or more amino acid residues introduced into it from a source which is non-human. These non-human amino acid residues are often referred to as "import" residues, which are typically taken from an "import" variable domain. Humanization can be essentially performed following the method of Winter and co-workers (Jones *et al.*, *Nature*, 321:522-525 (1986); Riechmann *et al.*, *Nature*, 332:323-327 (1988); Verhoeyen *et al.*, *Science*, 239:1534-1536 (1988)), by substituting rodent CDRs or CDR sequences for the corresponding sequences of a human antibody. Accordingly, such "humanized" antibodies are chimeric antibodies (U.S. Patent No. 4,816,567) wherein substantially less than an intact human variable domain has been substituted by the corresponding sequence from a non-human species. In practice, humanized antibodies are typically human antibodies in which some CDR residues and possibly some FR residues are substituted by residues from analogous sites in rodent antibodies.

The choice of human variable domains, both light and heavy, to be used in making the humanized antibodies is very important to reduce antigenicity. According to the so-called "best-fit" method, the sequence of the variable domain of a rodent antibody is screened against the entire library of known human variable-domain sequences. The human sequence which is closest to that of the rodent is then accepted as the human framework (FR) for the humanized antibody (Sims *et al.*, *J. Immunol.*, 151:2296 (1993); Chothia *et al.*, *J. Mol. Biol.*, 196:901 (1987)). Another method uses a particular framework derived from the consensus sequence of all human antibodies of a particular subgroup of light or heavy chains. The same framework may be used for several different humanized antibodies (Carter *et al.*, *Proc. Natl. Acad. Sci. USA*, 89:4285 (1992); Presta *et al.*, *J. Immunol.*, 151:2623 (1993)).

It is further important that antibodies be humanized with retention of high affinity for the antigen and other favorable biological properties. To achieve this goal, according to a preferred method, humanized antibodies are prepared by a process of analysis of the parental sequences and various conceptual humanized products using three-dimensional models of the parental and humanized sequences. Three-dimensional immunoglobulin models are commonly available and are familiar to those skilled in the art. Computer programs are available which illustrate and display probable three-dimensional conformational structures of selected candidate immunoglobulin sequences. Inspection of these displays permits analysis of the likely role of the residues in the functioning of the candidate immunoglobulin sequence, *i.e.*, the analysis of residues that influence the ability of the candidate immunoglobulin to bind its antigen. In this way, FR residues can be selected and combined from the recipient and import sequences so that the desired antibody characteristic, such as increased affinity for the target antigen(s), is achieved.

In general, the CDR residues are directly and most substantially involved in influencing antigen binding.

Alternatively, it is now possible to produce transgenic animals (e.g., mice) that are capable, upon immunization, of producing a full repertoire of human antibodies in the absence of endogenous immunoglobulin production. For example, it has been described that the homozygous deletion of the antibody heavy-chain joining region (J_H) gene in chimeric and germ-line mutant mice results in complete inhibition of endogenous antibody production. Transfer of the human germ-line immunoglobulin gene array in such germ-line mutant mice will result in the production of human antibodies upon antigen challenge. See, e.g., Jakobovits *et al.*, *Proc. Natl. Acad. Sci. USA*, 90:2551 (1993); Jakobovits *et al.*, *Nature*, 362:255-258 (1993); Bruggemann *et al.*, *Year in Immuno.*, 7:33 (1993); and Duchosal *et al.*, *Nature* 355:258 (1992). Human antibodies can also be derived from phage-display libraries (Hoogenboom *et al.*, *J. Mol. Biol.*, 227:381 (1991); Marks *et al.*, *J. Mol. Biol.*, 222:581-597 (1991); Vaughan *et al.*, *Nature Biotech* 14:309 (1996)).

15 (v) Multispecific antibodies

Multispecific antibodies have binding specificities for at least two different antigens. While such molecules normally will only bind two antigens (*i.e.* bispecific antibodies, BsAbs), antibodies with additional specificities such as trispecific antibodies are encompassed by this expression when used herein. Examples of BsAbs include those with one arm directed against a tumor cell antigen and the other arm directed against a cytotoxic trigger molecule such as anti-Fc γ RI/anti-CD15, anti-p185^{HER2}/Fc γ RIII (CD16), anti-CD3/anti-malignant B-cell (1D10), anti-CD3/anti-p185^{HER2}, anti-CD3/anti-p97, anti-CD3/anti-renal cell carcinoma, anti-CD3/anti-OVCAR-3, anti-CD3/L-D1 (anti-colon carcinoma), anti-CD3/anti-melanocyte stimulating hormone analog, anti-EGF receptor/anti-CD3, anti-CD3/anti-CAMA1, anti-CD3/anti-CD19, anti-CD3/MoV18, anti-neural cell adhesion molecule (NCAM)/anti-CD3, anti-folate binding protein (FBP)/anti-CD3, anti-pan carcinoma associated antigen (AMOC-31)/anti-CD3; BsAbs with one arm which binds specifically to a tumor antigen and one arm which binds to a toxin such as anti-saporin/anti-Id-1, anti-CD22/anti-saporin, anti-CD7/anti-saporin, anti-CD38/anti-saporin, anti-CEA/anti-ricin A chain, anti-interferon- α (IFN- α)/anti-hybridoma idiotype, anti-CEA/anti-vinca alkaloid; BsAbs for converting enzyme activated prodrugs such as anti-CD30/anti-alkaline phosphatase (which catalyzes conversion of mitomycin phosphate prodrug to mitomycin alcohol); BsAbs which can be used as fibrinolytic agents such as anti-fibrin/anti-tissue plasminogen activator (tPA), anti-fibrin/anti-urokinase-type plasminogen activator (uPA); BsAbs for targeting immune complexes to cell surface receptors such as anti-low density lipoprotein (LDL)/anti-Fc receptor (e.g. Fc γ RI, Fc γ RII or Fc γ RIII); BsAbs for use in therapy of infectious diseases such as anti-CD3/anti-herpes simplex

virus (HSV), anti-T-cell receptor:CD3 complex/anti-influenza, anti-FcγR/anti-HIV; BsAbs for tumor detection *in vitro* or *in vivo* such as anti-CEA/anti-EOTUBE, anti-CEA/anti-DPTA, anti-p185^{HER2}/anti-hapten; BsAbs as vaccine adjuvants; and BsAbs as diagnostic tools such as anti-rabbit IgG/anti-ferritin, anti-horse radish peroxidase (HRP)/anti-hormone, anti-somatostatin/anti-substance P, anti-HRP/anti-FITC, anti-CEA/anti-β-galactosidase. Examples of trispecific antibodies include anti-CD3/anti-CD4/anti-CD37, anti-CD3/anti-CD5/anti-CD37 and anti-CD3/anti-CD8/anti-CD37. Bispecific antibodies can be prepared as full length antibodies or antibody fragments (e.g. F(ab')₂ bispecific antibodies).

Methods for making bispecific antibodies are known in the art. Traditional production of full length bispecific antibodies is based on the coexpression of two immunoglobulin heavy chain-light chain pairs, where the two chains have different specificities (Millstein *et al.*, *Nature*, 305:537-539 (1983)). Because of the random assortment of immunoglobulin heavy and light chains, these hybridomas (quadromas) produce a potential mixture of 10 different antibody molecules, of which only one has the correct bispecific structure. Purification of the correct molecule, which is usually done by affinity chromatography steps, is rather cumbersome, and the product yields are low. Similar procedures are disclosed in WO 93/08829, and in Traunecker *et al.*, *EMBO J.*, 10:3655-3659 (1991).

According to a different approach, antibody variable domains with the desired binding specificities (antibody-antigen combining sites) are fused to immunoglobulin constant domain sequences. The fusion preferably is with an immunoglobulin heavy chain constant domain, comprising at least part of the hinge, CH2, and CH3 regions. It is preferred to have the first heavy-chain constant region (CH1) containing the site necessary for light chain binding, present in at least one of the fusions. DNAs encoding the immunoglobulin heavy chain fusions and, if desired, the immunoglobulin light chain, are inserted into separate expression vectors, and are co-transfected into a suitable host organism. This provides for great flexibility in adjusting the mutual proportions of the three polypeptide fragments in embodiments when unequal ratios of the three polypeptide chains used in the construction provide the optimum yields. It is, however, possible to insert the coding sequences for two or all three polypeptide chains in one expression vector when the expression of at least two polypeptide chains in equal ratios results in high yields or when the ratios are of no particular significance.

In a preferred embodiment of this approach, the bispecific antibodies are composed of a hybrid immunoglobulin heavy chain with a first binding specificity in one arm, and a hybrid immunoglobulin heavy chain-light chain pair (providing a second binding specificity) in the other arm. It was found that this asymmetric structure facilitates the separation of the desired bispecific compound from unwanted immunoglobulin chain combinations, as the presence of an

immunoglobulin light chain in only one half of the bispecific molecule provides for a facile way of separation. This approach is disclosed in WO 94/04690. For further details of generating bispecific antibodies see, for example, Suresh *et al.*, *Methods in Enzymology*, 121:210 (1986). According to another approach described in W096/27011, the interface between a pair of antibody
5 molecules can be engineered to maximize the percentage of heterodimers which are recovered from recombinant cell culture. The preferred interface comprises at least a part of the C_H3 domain of an antibody constant domain. In this method, one or more small amino acid side chains from the interface of the first antibody molecule are replaced with larger side chains (*e.g.* tyrosine or tryptophan). Compensatory "cavities" of identical or similar size to the large side chain(s) are
10 created on the interface of the second antibody molecule by replacing large amino acid side chains with smaller ones (*e.g.* alanine or threonine). This provides a mechanism for increasing the yield of the heterodimer over other unwanted end-products such as homodimers.

Bispecific antibodies include cross-linked or "heteroconjugate" antibodies. For example, one of the antibodies in the heteroconjugate can be coupled to avidin, the other to biotin. Such
15 antibodies have, for example, been proposed to target immune system cells to unwanted cells (US Patent No. 4,676,980), and for treatment of HIV infection (WO 91/00360, WO 92/200373, and EP 03089). Heteroconjugate antibodies may be made using any convenient cross-linking methods. Suitable cross-linking agents are well known in the art, and are disclosed in US Patent No. 4,676,980, along with a number of cross-linking techniques.

20 Antibodies with more than two valencies are contemplated. For example, trispecific antibodies can be prepared. Tutt *et al.* *J. Immunol.* 147: 60 (1991).

While the polypeptide of interest herein is preferably an antibody, other Fc region-containing polypeptides which can be modified according to the methods described herein are contemplated. An example of such a molecule is an immunoadhesin.

25 C. Immunoadhesin Preparation

The simplest and most straightforward immunoadhesin design combines the binding domain(s) of the adhesin (*e.g.* the extracellular domain (ECD) of a receptor) with the Fc region of an immunoglobulin heavy chain. Ordinarily, when preparing the immunoadhesins of the present invention, nucleic acid encoding the binding domain of the adhesin will be fused C-
30 terminally to nucleic acid encoding the N-terminus of an immunoglobulin constant domain sequence, however N-terminal fusions are also possible.

Typically, in such fusions the encoded chimeric polypeptide will retain at least functionally active hinge, C_H2 and C_H3 domains of the constant region of an immunoglobulin heavy chain. Fusions are also made to the C-terminus of the Fc portion of a constant domain, or immediately
35 N-terminal to the C_H1 of the heavy chain or the corresponding region of the light chain. The

precise site at which the fusion is made is not critical; particular sites are well known and may be selected in order to optimize the biological activity, secretion, or binding characteristics of the immunoadhesin.

In a preferred embodiment, the adhesin sequence is fused to the N-terminus of the Fc region of immunoglobulin G₁ (IgG₁). It is possible to fuse the entire heavy chain constant region to the adhesin sequence. However, more preferably, a sequence beginning in the hinge region just upstream of the papain cleavage site which defines IgG Fc chemically (*i.e.* residue 216, taking the first residue of heavy chain constant region to be 114), or analogous sites of other immunoglobulins is used in the fusion. In a particularly preferred embodiment, the adhesin amino acid sequence is fused to (a) the hinge region and C_H2 and C_H3 or (b) the C_H1, hinge, C_H2 and C_H3 domains, of an IgG heavy chain.

For bispecific immunoadhesins, the immunoadhesins are assembled as multimers, and particularly as heterodimers or heterotetramers. Generally, these assembled immunoglobulins will have known unit structures. A basic four chain structural unit is the form in which IgG, IgD, and IgE exist. A four chain unit is repeated in the higher molecular weight immunoglobulins; IgM generally exists as a pentamer of four basic units held together by disulfide bonds. IgA globulin, and occasionally IgG globulin, may also exist in multimeric form in serum. In the case of multimer, each of the four units may be the same or different.

Various exemplary assembled immunoadhesins within the scope herein are schematically diagrammed below:

- (a) AC_L-AC_L;
- (b) AC_H-(AC_H, AC_L-AC_H, AC_L-V_HC_H, or V_LC_L-AC_H);
- (c) AC_L-AC_H-(AC_L-AC_H, AC_L-V_HC_H, V_LC_L-AC_H, or V_LC_L-V_HC_H);
- (d) AC_L-V_HC_H-(AC_H, or AC_L-V_HC_H, or V_LC_L-AC_H);
- (e) V_LC_L-AC_H-(AC_L-V_HC_H, or V_LC_L-AC_H); and
- (f) (A-Y)_n-(V_LC_L-V_HC_H)₂,

wherein each A represents identical or different adhesin amino acid sequences;

V_L is an immunoglobulin light chain variable domain;

V_H is an immunoglobulin heavy chain variable domain;

C_L is an immunoglobulin light chain constant domain;

C_H is an immunoglobulin heavy chain constant domain;

n is an integer greater than 1;

Y designates the residue of a covalent cross-linking agent.

In the interests of brevity, the foregoing structures only show key features; they do not indicate joining (J) or other domains of the immunoglobulins, nor are disulfide bonds shown.

However, where such domains are required for binding activity, they shall be constructed to be present in the ordinary locations which they occupy in the immunoglobulin molecules.

Alternatively, the adhesin sequences can be inserted between immunoglobulin heavy chain and light chain sequences, such that an immunoglobulin comprising a chimeric heavy chain is obtained. In this embodiment, the adhesin sequences are fused to the 3' end of an immunoglobulin heavy chain in each arm of an immunoglobulin, either between the hinge and the C_H2 domain, or between the C_H2 and C_H3 domains. Similar constructs have been reported by Hoogenboom, *et al.*, *Mol. Immunol.* 28:1027-1037 (1991).

Although the presence of an immunoglobulin light chain is not required in the immunoadhesins of the present invention, an immunoglobulin light chain might be present either covalently associated to an adhesin-immunoglobulin heavy chain fusion polypeptide, or directly fused to the adhesin. In the former case, DNA encoding an immunoglobulin light chain is typically coexpressed with the DNA encoding the adhesin-immunoglobulin heavy chain fusion protein. Upon secretion, the hybrid heavy chain and the light chain will be covalently associated to provide an immunoglobulin-like structure comprising two disulfide-linked immunoglobulin heavy chain-light chain pairs. Methods suitable for the preparation of such structures are, for example, disclosed in U.S. Patent No. 4,816,567, issued 28 March 1989.

Immunoadhesins are most conveniently constructed by fusing the cDNA sequence encoding the adhesin portion in-frame to an immunoglobulin cDNA sequence. However, fusion to genomic immunoglobulin fragments can also be used (see, *e.g.* Aruffo *et al.*, *Cell* 61:1303-1313 (1990); and Stamenkovic *et al.*, *Cell* 66:1133-1144 (1991)). The latter type of fusion requires the presence of Ig regulatory sequences for expression. cDNAs encoding IgG heavy-chain constant regions can be isolated based on published sequences from cDNA libraries derived from spleen or peripheral blood lymphocytes, by hybridization or by polymerase chain reaction (PCR) techniques. The cDNAs encoding the "adhesin" and the immunoglobulin parts of the immunoadhesin are inserted in tandem into a plasmid vector that directs efficient expression in the chosen host cells.

D. Vectors, Host Cells and Recombinant Methods

The invention also provides isolated nucleic acid encoding a polypeptide variant as disclosed herein, vectors and host cells comprising the nucleic acid, and recombinant techniques for the production of the polypeptide variant.

For recombinant production of the polypeptide variant, the nucleic acid encoding it is isolated and inserted into a replicable vector for further cloning (amplification of the DNA) or for expression. DNA encoding the polypeptide variant is readily isolated and sequenced using conventional procedures (*e.g.*, by using oligonucleotide probes that are capable of binding

specifically to genes encoding the polypeptide variant). Many vectors are available. The vector components generally include, but are not limited to, one or more of the following: a signal sequence, an origin of replication, one or more marker genes, an enhancer element, a promoter, and a transcription termination sequence.

5 (i) **Signal sequence component**

The polypeptide variant of this invention may be produced recombinantly not only directly, but also as a fusion polypeptide with a heterologous polypeptide, which is preferably a signal sequence or other polypeptide having a specific cleavage site at the N-terminus of the mature protein or polypeptide. The heterologous signal sequence selected preferably is one that is
10 recognized and processed (*i.e.*, cleaved by a signal peptidase) by the host cell. For prokaryotic host cells that do not recognize and process the native polypeptide variant signal sequence, the signal sequence is substituted by a prokaryotic signal sequence selected, for example, from the group of the alkaline phosphatase, penicillinase, lpp, or heat-stable enterotoxin II leaders. For yeast secretion the native signal sequence may be substituted by, *e.g.*, the yeast invertase leader,
15 α factor leader (including *Saccharomyces* and *Kluyveromyces* α -factor leaders), or acid phosphatase leader, the *C. albicans* glucoamylase leader, or the signal described in WO 90/13646. In mammalian cell expression, mammalian signal sequences as well as viral secretory leaders, for example, the herpes simplex gD signal, are available.

The DNA for such precursor region is ligated in reading frame to DNA encoding the
20 polypeptide variant.

 (ii) **Origin of replication component**

Both expression and cloning vectors contain a nucleic acid sequence that enables the vector to replicate in one or more selected host cells. Generally, in cloning vectors this sequence is one that enables the vector to replicate independently of the host chromosomal DNA, and
25 includes origins of replication or autonomously replicating sequences. Such sequences are well known for a variety of bacteria, yeast, and viruses. The origin of replication from the plasmid pBR322 is suitable for most Gram-negative bacteria, the 2 μ plasmid origin is suitable for yeast, and various viral origins (SV40, polyoma, adenovirus, VSV or BPV) are useful for cloning vectors in mammalian cells. Generally, the origin of replication component is not needed for mammalian
30 expression vectors (the SV40 origin may typically be used only because it contains the early promoter).

 (iii) **Selection gene component**

Expression and cloning vectors may contain a selection gene, also termed a selectable marker. Typical selection genes encode proteins that (a) confer resistance to antibiotics or other
35 toxins, *e.g.*, ampicillin, neomycin, methotrexate, or tetracycline, (b) complement auxotrophic

deficiencies, or (c) supply critical nutrients not available from complex media, *e.g.*, the gene encoding D-alanine racemase for *Bacillus*.

One example of a selection scheme utilizes a drug to arrest growth of a host cell. Those cells that are successfully transformed with a heterologous gene produce a protein conferring drug resistance and thus survive the selection regimen. Examples of such dominant selection use the drugs neomycin, mycophenolic acid and hygromycin.

Another example of suitable selectable markers for mammalian cells are those that enable the identification of cells competent to take up the polypeptide variant nucleic acid, such as DHFR, thymidine kinase, metallothionein-I and -II, preferably primate metallothionein genes, adenosine deaminase, ornithine decarboxylase, *etc.*

For example, cells transformed with the DHFR selection gene are first identified by culturing all of the transformants in a culture medium that contains methotrexate (Mtx), a competitive antagonist of DHFR. An appropriate host cell when wild-type DHFR is employed is the Chinese hamster ovary (CHO) cell line deficient in DHFR activity.

Alternatively, host cells (particularly wild-type hosts that contain endogenous DHFR) transformed or co-transformed with DNA sequences encoding polypeptide variant, wild-type DHFR protein, and another selectable marker such as aminoglycoside 3'-phosphotransferase (APH) can be selected by cell growth in medium containing a selection agent for the selectable marker such as an aminoglycosidic antibiotic, *e.g.*, kanamycin, neomycin, or G418. See U.S. Patent No. 4,965,199.

A suitable selection gene for use in yeast is the *trp1* gene present in the yeast plasmid YRp7 (Stinchcomb *et al.*, *Nature*, 282:39 (1979)). The *trp1* gene provides a selection marker for a mutant strain of yeast lacking the ability to grow in tryptophan, for example, ATCC No. 44076 or PEP4-1. Jones, *Genetics*, 85:12 (1977). The presence of the *trp1* lesion in the yeast host cell genome then provides an effective environment for detecting transformation by growth in the absence of tryptophan. Similarly, *Leu2*-deficient yeast strains (ATCC 20,622 or 38,626) are complemented by known plasmids bearing the *Leu2* gene.

In addition, vectors derived from the 1.6 μ m circular plasmid pKD1 can be used for transformation of *Kluyveromyces* yeasts. Alternatively, an expression system for large-scale production of recombinant calf chymosin was reported for *K. lactis*. Van den Berg, *Bio/Technology*, 8:135 (1990). Stable multi-copy expression vectors for secretion of mature recombinant human serum albumin by industrial strains of *Kluyveromyces* have also been disclosed. Fleer *et al.*, *Bio/Technology*, 9:968-975 (1991).

(iv) **Promoter component**

Expression and cloning vectors usually contain a promoter that is recognized by the host organism and is operably linked to the polypeptide variant nucleic acid. Promoters suitable for use with prokaryotic hosts include the *phoA* promoter, β -lactamase and lactose promoter systems, alkaline phosphatase, a tryptophan (*trp*) promoter system, and hybrid promoters such as the *tac* promoter. However, other known bacterial promoters are suitable. Promoters for use in bacterial systems also will contain a Shine-Dalgarno (S.D.) sequence operably linked to the DNA encoding the polypeptide variant.

Promoter sequences are known for eukaryotes. Virtually all eukaryotic genes have an AT-rich region located approximately 25 to 30 bases upstream from the site where transcription is initiated. Another sequence found 70 to 80 bases upstream from the start of transcription of many genes is a CNCAAT region where N may be any nucleotide. At the 3' end of most eukaryotic genes is an AATAAA sequence that may be the signal for addition of the poly A tail to the 3' end of the coding sequence. All of these sequences are suitably inserted into eukaryotic expression vectors.

Examples of suitable promoting sequences for use with yeast hosts include the promoters for 3-phosphoglycerate kinase or other glycolytic enzymes, such as enolase, glyceraldehyde-3-phosphate dehydrogenase, hexokinase, pyruvate decarboxylase, phosphofructokinase, glucose-6-phosphate isomerase, 3-phosphoglycerate mutase, pyruvate kinase, triosephosphate isomerase, phosphoglucose isomerase, and glucokinase.

Other yeast promoters, which are inducible promoters having the additional advantage of transcription controlled by growth conditions, are the promoter regions for alcohol dehydrogenase 2, isocytocrome C, acid phosphatase, degradative enzymes associated with nitrogen metabolism, metallothionein, glyceraldehyde-3-phosphate dehydrogenase, and enzymes responsible for maltose and galactose utilization. Suitable vectors and promoters for use in yeast expression are further described in EP 73,657. Yeast enhancers also are advantageously used with yeast promoters.

Polypeptide variant transcription from vectors in mammalian host cells is controlled, for example, by promoters obtained from the genomes of viruses such as polyoma virus, fowlpox virus, adenovirus (such as Adenovirus 2), bovine papilloma virus, avian sarcoma virus, cytomegalovirus, a retrovirus, hepatitis-B virus and most preferably Simian Virus 40 (SV40), from heterologous mammalian promoters, *e.g.*, the actin promoter or an immunoglobulin promoter, from heat-shock promoters, provided such promoters are compatible with the host cell systems.

The early and late promoters of the SV40 virus are conveniently obtained as an SV40 restriction fragment that also contains the SV40 viral origin of replication. The immediate early

promoter of the human cytomegalovirus is conveniently obtained as a HindIII E restriction fragment. A system for expressing DNA in mammalian hosts using the bovine papilloma virus as a vector is disclosed in U.S. Patent No. 4,419,446. A modification of this system is described in U.S. Patent No. 4,601,978. See also Reyes *et al.*, *Nature* 297:598-601 (1982) on expression of
5 human β -interferon cDNA in mouse cells under the control of a thymidine kinase promoter from herpes simplex virus. Alternatively, the rous sarcoma virus long terminal repeat can be used as the promoter.

(v) Enhancer element component

Transcription of a DNA encoding the polypeptide variant of this invention by higher
10 eukaryotes is often increased by inserting an enhancer sequence into the vector. Many enhancer sequences are now known from mammalian genes (globin, elastase, albumin, α -fetoprotein, and insulin). Typically, however, one will use an enhancer from a eukaryotic cell virus. Examples include the SV40 enhancer on the late side of the replication origin (bp 100-270), the cytomegalovirus early promoter enhancer, the polyoma enhancer on the late side of the
15 replication origin, and adenovirus enhancers. See also Yaniv, *Nature* 297:17-18 (1982) on enhancing elements for activation of eukaryotic promoters. The enhancer may be spliced into the vector at a position 5' or 3' to the polypeptide variant-encoding sequence, but is preferably located at a site 5' from the promoter.

(vi) Transcription termination component

20 Expression vectors used in eukaryotic host cells (yeast, fungi, insect, plant, animal, human, or nucleated cells from other multicellular organisms) will also contain sequences necessary for the termination of transcription and for stabilizing the mRNA. Such sequences are commonly available from the 5' and, occasionally 3', untranslated regions of eukaryotic or viral DNAs or cDNAs. These regions contain nucleotide segments transcribed as polyadenylated
25 fragments in the untranslated portion of the mRNA encoding the polypeptide variant. One useful transcription termination component is the bovine growth hormone polyadenylation region. See WO94/11026 and the expression vector disclosed therein.

(vii) Selection and transformation of host cells

Suitable host cells for cloning or expressing the DNA in the vectors herein are the
30 prokaryote, yeast, or higher eukaryote cells described above. Suitable prokaryotes for this purpose include eubacteria, such as Gram-negative or Gram-positive organisms, for example, Enterobacteriaceae such as *Escherichia*, e.g., *E. coli*, *Enterobacter*, *Erwinia*, *Klebsiella*, *Proteus*, *Salmonella*, e.g., *Salmonella typhimurium*, *Serratia*, e.g., *Serratia marcescans*, and *Shigella*, as well as *Bacilli* such as *B. subtilis* and *B. licheniformis* (e.g., *B. licheniformis* 41P
35 disclosed in DD 266,710 published 12 April 1989), *Pseudomonas* such as *P. aeruginosa*, and

Streptomyces. One preferred *E. coli* cloning host is *E. coli* 294 (ATCC 31,446), although other strains such as *E. coli* B, *E. coli* X1776 (ATCC 31,537), and *E. coli* W3110 (ATCC 27,325) are suitable. These examples are illustrative rather than limiting.

In addition to prokaryotes, eukaryotic microbes such as filamentous fungi or yeast are suitable cloning or expression hosts for polypeptide variant-encoding vectors. **Saccharomyces cerevisiae**, or common baker's yeast, is the most commonly used among lower eukaryotic host microorganisms. However, a number of other genera, species, and strains are commonly available and useful herein, such as **Schizosaccharomyces pombe**; **Kluyveromyces** hosts such as, e.g., *K. lactis*, *K. fragilis* (ATCC 12,424), *K. bulgaricus* (ATCC 16,045), *K. wickerhamii* (ATCC 24,178), *K. waltii* (ATCC 56,500), *K. drosophilae* (ATCC 36,906), *K. thermotolerans*, and *K. marxianus*; **Yarrowia** (EP 402,226); **Pichia pastoris** (EP 183,070); **Candida**; **Trichoderma reesei** (EP 244,234); **Neurospora crassa**; **Schwanniomyces** such as **Schwanniomyces occidentalis**; and filamentous fungi such as, e.g., **Neurospora**, **Penicillium**, **Tolypocladium**, and **Aspergillus** hosts such as *A. nidulans* and *A. niger*.

Suitable host cells for the expression of glycosylated polypeptide variant are derived from multicellular organisms. Examples of invertebrate cells include plant and insect cells. Numerous baculoviral strains and variants and corresponding permissive insect host cells from hosts such as **Spodoptera frugiperda** (caterpillar), **Aedes aegypti** (mosquito), **Aedes albopictus** (mosquito), **Drosophila melanogaster** (fruitfly), and **Bombyx mori** have been identified. A variety of viral strains for transfection are publicly available, e.g., the L-1 variant of **Autographa californica** NPV and the Bm-5 strain of **Bombyx mori** NPV, and such viruses may be used as the virus herein according to the present invention, particularly for transfection of **Spodoptera frugiperda** cells.

Plant cell cultures of cotton, corn, potato, soybean, petunia, tomato, and tobacco can also be utilized as hosts.

However, interest has been greatest in vertebrate cells, and propagation of vertebrate cells in culture (tissue culture) has become a routine procedure. Examples of useful mammalian host cell lines are monkey kidney CV1 line transformed by SV40 (COS-7, ATCC CRL 1651); human embryonic kidney line (293 or 293 cells subcloned for growth in suspension culture, Graham *et al.*, *J. Gen Virol.* 36:59 (1977)); baby hamster kidney cells (BHK, ATCC CCL 10); Chinese hamster ovary cells/-DHFR (CHO, Urlaub *et al.*, *Proc. Natl. Acad. Sci. USA* 77:4216 (1980)); mouse sertoli cells (TM4, Mather, *Biol. Reprod.* 23:243-251 (1980)); monkey kidney cells (CV1 ATCC CCL 70); African green monkey kidney cells (VERO-76, ATCC CRL-1587); human cervical carcinoma cells (HELA, ATCC CCL 2); canine kidney cells (MDCK, ATCC CCL 34); buffalo rat liver cells (BRL 3A, ATCC CRL 1442); human lung cells (W138, ATCC CCL 75); human

liver cells (Hep G2, HB 8065); mouse mammary tumor (MMT 060562, ATCC CCL51); TRI cells (Mather *et al.*, *Annals N.Y. Acad. Sci.* 383:44-68 (1982)); MRC 5 cells; FS4 cells; and a human hepatoma line (Hep G2).

5 Host cells are transformed with the above-described expression or cloning vectors for polypeptide variant production and cultured in conventional nutrient media modified as appropriate for inducing promoters, selecting transformants, or amplifying the genes encoding the desired sequences.

(viii) Culturing the host cells

The host cells used to produce the polypeptide variant of this invention may be cultured
10 in a variety of media. Commercially available media such as Ham's F10 (Sigma), Minimal Essential Medium ((MEM), (Sigma), RPMI-1640 (Sigma), and Dulbecco's Modified Eagle's Medium ((DMEM), Sigma) are suitable for culturing the host cells. In addition, any of the media described in Ham *et al.*, *Meth. Enz.* 58:44 (1979), Barnes *et al.*, *Anal. Biochem.* 102:255 (1980), U.S. Pat. Nos. 4,767,704; 4,657,866; 4,927,762; 4,560,655; or 5,122,469; WO 90/03430; WO
15 87/00195; or U.S. Patent Re. 30,985 may be used as culture media for the host cells. Any of these media may be supplemented as necessary with hormones and/or other growth factors (such as insulin, transferrin, or epidermal growth factor), salts (such as sodium chloride, calcium, magnesium, and phosphate), buffers (such as HEPES), nucleotides (such as adenosine and thymidine), antibiotics (such as GENTAMYCIN™ drug), trace elements (defined as inorganic
20 compounds usually present at final concentrations in the micromolar range), and glucose or an equivalent energy source. Any other necessary supplements may also be included at appropriate concentrations that would be known to those skilled in the art. The culture conditions, such as temperature, pH, and the like, are those previously used with the host cell selected for expression, and will be apparent to the ordinarily skilled artisan.

25 **(ix) Polypeptide variant purification**

When using recombinant techniques, the polypeptide variant can be produced intracellularly, in the periplasmic space, or directly secreted into the medium. If the polypeptide variant is produced intracellularly, as a first step, the particulate debris, either host cells or lysed fragments, is removed, for example, by centrifugation or ultrafiltration. Carter *et al.*,
30 *Bio/Technology* 10:163-167 (1992) describe a procedure for isolating antibodies which are secreted to the periplasmic space of *E. coli*. Briefly, cell paste is thawed in the presence of sodium acetate (pH 3.5), EDTA, and phenylmethylsulfonylfluoride (PMSF) over about 30 min. Cell debris can be removed by centrifugation. Where the polypeptide variant is secreted into the medium, supernatants from such expression systems are generally first concentrated using a
35 commercially available protein concentration filter, for example, an Amicon or Millipore Pellicon

ultrafiltration unit. A protease inhibitor such as PMSF may be included in any of the foregoing steps to inhibit proteolysis and antibiotics may be included to prevent the growth of adventitious contaminants.

5 The polypeptide variant composition prepared from the cells can be purified using, for example, hydroxylapatite chromatography, gel electrophoresis, dialysis, and affinity chromatography, with affinity chromatography being the preferred purification technique. The suitability of protein A as an affinity ligand depends on the species and isotype of any immunoglobulin Fc region that is present in the polypeptide variant. Protein A can be used to purify polypeptide variants that are based on human $\gamma 1$, $\gamma 2$, or $\gamma 4$ heavy chains (Lindmark *et al.*,
10 *J. Immunol. Meth.* 62:1-13 (1983)). Protein G is recommended for all mouse isotypes and for human $\gamma 3$ (Guss *et al.*, *EMBO J.* 5:15671575 (1986)). The matrix to which the affinity ligand is attached is most often agarose, but other matrices are available. Mechanically stable matrices such as controlled pore glass or poly(styrenedivinyl)benzene allow for faster flow rates and shorter processing times than can be achieved with agarose. Where the polypeptide variant
15 comprises a C_H3 domain, the Bakerbond ABX™ resin (J. T. Baker, Phillipsburg, NJ) is useful for purification. Other techniques for protein purification such as fractionation on an ion-exchange column, ethanol precipitation, Reverse Phase HPLC, chromatography on silica, chromatography on heparin SEPHAROSE™ chromatography on an anion or cation exchange resin (such as a polyaspartic acid column), chromatofocusing, SDS-PAGE, and ammonium sulfate precipitation
20 are also available depending on the polypeptide variant to be recovered.

Following any preliminary purification step(s), the mixture comprising the polypeptide variant of interest and contaminants may be subjected to low pH hydrophobic interaction chromatography using an elution buffer at a pH between about 2.5-4.5, preferably performed at low salt concentrations (*e.g.*, from about 0-0.25M salt).

25 E. Pharmaceutical Formulations

Therapeutic formulations of the polypeptide variant are prepared for storage by mixing the polypeptide variant having the desired degree of purity with optional physiologically acceptable carriers, excipients or stabilizers (*Remington's Pharmaceutical Sciences* 16th edition, Osol, A. Ed. (1980)), in the form of lyophilized formulations or aqueous solutions. Acceptable carriers,
30 excipients, or stabilizers are nontoxic to recipients at the dosages and concentrations employed, and include buffers such as phosphate, citrate, and other organic acids; antioxidants including ascorbic acid and methionine; preservatives (such as octadecyldimethylbenzyl ammonium chloride; hexamethonium chloride; benzalkonium chloride, benzethonium chloride; phenol, butyl or benzyl alcohol; alkyl parabens such as methyl or propyl paraben; catechol; resorcinol;
35 cyclohexanol; 3-pentanol; and m-cresol); low molecular weight (less than about 10 residues)

polypeptide; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, histidine, arginine, or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrans; chelating agents such as EDTA; sugars such as sucrose, mannitol, trehalose or sorbitol; salt-forming counter-ions such as sodium; metal complexes (e.g., Zn-protein complexes); and/or non-ionic surfactants such as TWEEN™, PLURONICS™ or polyethylene glycol (PEG).

The formulation herein may also contain more than one active compound as necessary for the particular indication being treated, preferably those with complementary activities that do not adversely affect each other. Such molecules are suitably present in combination in amounts that are effective for the purpose intended.

The active ingredients may also be entrapped in microcapsule prepared, for example, by coacervation techniques or by interfacial polymerization, for example, hydroxymethylcellulose or gelatin-microcapsule and poly-(methylmethacrylate) microcapsule, respectively, in colloidal drug delivery systems (for example, liposomes, albumin microspheres, microemulsions, nano-particles and nanocapsules) or in macroemulsions. Such techniques are disclosed in *Remington's Pharmaceutical Sciences* 16th edition, Osol, A. Ed. (1980).

The formulations to be used for *in vivo* administration must be sterile. This is readily accomplished by filtration through sterile filtration membranes.

Sustained-release preparations may be prepared. Suitable examples of sustained-release preparations include semipermeable matrices of solid hydrophobic polymers containing the polypeptide variant, which matrices are in the form of shaped articles, e.g., films, or microcapsule. Examples of sustained-release matrices include polyesters, hydrogels (for example, poly(2-hydroxyethyl-methacrylate), or poly(vinylalcohol)), polylactides (U.S. Pat. No. 3,773,919), copolymers of L-glutamic acid and γ ethyl-L-glutamate, non-degradable ethylene-vinyl acetate, degradable lactic acid-glycolic acid copolymers such as the LUPRON DEPOT™ (injectable microspheres composed of lactic acid-glycolic acid copolymer and leuprolide acetate), and poly-D-(-)-3-hydroxybutyric acid. While polymers such as ethylene-vinyl acetate and lactic acid-glycolic acid enable release of molecules for over 100 days, certain hydrogels release proteins for shorter time periods. When encapsulated antibodies remain in the body for a long time, they may denature or aggregate as a result of exposure to moisture at 37°C, resulting in a loss of biological activity and possible changes in immunogenicity. Rational strategies can be devised for stabilization depending on the mechanism involved. For example, if the aggregation mechanism is discovered to be intermolecular S-S bond formation through thio-disulfide interchange, stabilization may be achieved by modifying sulfhydryl residues, lyophilizing from

acidic solutions, controlling moisture content, using appropriate additives, and developing specific polymer matrix compositions.

F. Non-Therapeutic Uses for the Polypeptide Variant

The polypeptide variant of the invention may be used as an affinity purification agent. In this process, the polypeptide variant is immobilized on a solid phase such as a Sephadex resin or filter paper, using methods well known in the art. The immobilized polypeptide variant is contacted with a sample containing the antigen to be purified, and thereafter the support is washed with a suitable solvent that will remove substantially all the material in the sample except the antigen to be purified, which is bound to the immobilized polypeptide variant. Finally, the support is washed with another suitable solvent, such as glycine buffer, pH 5.0, that will release the antigen from the polypeptide variant.

The polypeptide variant may also be useful in diagnostic assays, *e.g.*, for detecting expression of an antigen of interest in specific cells, tissues, or serum.

For diagnostic applications, the polypeptide variant typically will be labeled with a detectable moiety. Numerous labels are available which can be generally grouped into the following categories:

(a) Radioisotopes, such as ^{35}S , ^{14}C , ^{125}I , ^3H , and ^{131}I . The polypeptide variant can be labeled with the radioisotope using the techniques described in *Current Protocols in Immunology*, Volumes 1 and 2, Coligen *et al.*, Ed. Wiley-Interscience, New York, New York, Pubs. (1991) for example and radioactivity can be measured using scintillation counting.

(b) Fluorescent labels such as rare earth chelates (europium chelates) or fluorescein and its derivatives, rhodamine and its derivatives, dansyl, Lissamine, phycoerythrin and Texas Red are available. The fluorescent labels can be conjugated to the polypeptide variant using the techniques disclosed in *Current Protocols in Immunology*, *supra*, for example. Fluorescence can be quantified using a fluorimeter.

(c) Various enzyme-substrate labels are available and U.S. Patent No. 4,275,149 provides a review of some of these. The enzyme generally catalyzes a chemical alteration of the chromogenic substrate that can be measured using various techniques. For example, the enzyme may catalyze a color change in a substrate, which can be measured spectrophotometrically. Alternatively, the enzyme may alter the fluorescence or chemiluminescence of the substrate. Techniques for quantifying a change in fluorescence are described above. The chemiluminescent substrate becomes electronically excited by a chemical reaction and may then emit light which can be measured (using a chemiluminometer, for example) or donates energy to a fluorescent acceptor. Examples of enzymatic labels include luciferases (*e.g.*, firefly luciferase and bacterial luciferase; U.S. Patent No. 4,737,456), luciferin, 2,3-dihydrophthalazinediones, malate

dehydrogenase, urease, peroxidase such as horseradish peroxidase (HRPO), alkaline phosphatase, β -galactosidase, glucoamylase, lysozyme, saccharide oxidases (e.g., glucose oxidase, galactose oxidase, and glucose-6-phosphate dehydrogenase), heterocyclic oxidases (such as uricase and xanthine oxidase), lactoperoxidase, microperoxidase, and the like.

- 5 Techniques for conjugating enzymes to antibodies are described in O'Sullivan *et al.*, Methods for the Preparation of Enzyme-Antibody Conjugates for use in Enzyme Immunoassay, in *Methods in Enzym.* (ed J. Langone & H. Van Vunakis), Academic press, New York, 73:147-166 (1981).

Examples of enzyme-substrate combinations include, for example:

- (i) Horseradish peroxidase (HRPO) with hydrogen peroxide as a substrate, wherein
10 the hydrogen peroxidase oxidizes a dye precursor (e.g., orthophenylene diamine (OPD) or 3,3',5,5'-tetramethyl benzidine hydrochloride (TMB));
- (ii) alkaline phosphatase (AP) with para-Nitrophenyl phosphate as chromogenic substrate; and
- (iii) β -D-galactosidase (β -D-Gal) with a chromogenic substrate (e.g., p-nitrophenyl- β -
15 D-galactosidase) or fluorogenic substrate 4-methylumbelliferyl- β -D-galactosidase.

Numerous other enzyme-substrate combinations are available to those skilled in the art. For a general review of these, see U.S. Patent Nos. 4,275,149 and 4,318,980.

- Sometimes, the label is indirectly conjugated with the polypeptide variant. The skilled artisan will be aware of various techniques for achieving this. For example, the polypeptide variant
20 can be conjugated with biotin and any of the three broad categories of labels mentioned above can be conjugated with avidin, or *vice versa*. Biotin binds selectively to avidin and thus, the label can be conjugated with the polypeptide variant in this indirect manner. Alternatively, to achieve indirect conjugation of the label with the polypeptide variant, the polypeptide variant is conjugated with a small hapten (e.g., digoxin) and one of the different types of labels mentioned above is
25 conjugated with an anti-hapten polypeptide variant (e.g., anti-digoxin antibody). Thus, indirect conjugation of the label with the polypeptide variant can be achieved.

In another embodiment of the invention, the polypeptide variant need not be labeled, and the presence thereof can be detected using a labeled antibody which binds to the polypeptide variant.

- 30 The polypeptide variant of the present invention may be employed in any known assay method, such as competitive binding assays, direct and indirect sandwich assays, and immunoprecipitation assays. Zola, *Monoclonal Antibodies: A Manual of Techniques*, pp.147-158 (CRC Press, Inc. 1987).

The polypeptide variant may also be used for *in vivo* diagnostic assays. Generally, the polypeptide variant is labeled with a radionuclide (such as ^{111}In , ^{99}Tc , ^{14}C , ^{131}I , ^{125}I , ^3H , ^{32}P or ^{35}S) so that the antigen or cells expressing it can be localized using immunoscintigraphy.

G. *In Vivo* Uses for the Polypeptide Variant

5 It is contemplated that the polypeptide variant of the present invention may be used to treat a mammal *e.g.* a patient suffering from, or predisposed to, a disease or disorder who could benefit from administration of the polypeptide variant. The conditions which can be treated with the polypeptide variant are many and include cancer (*e.g.* where the polypeptide variant binds the HER2 receptor, CD20 or vascular endothelial growth factor (VEGF)); allergic conditions such
10 as asthma (with an anti-IgE antibody); and LFA-1-mediated disorders (*e.g.* where the polypeptide variant is an anti-LFA-1 or anti-ICAM-1 antibody) etc.

Where the antibody binds the HER2 receptor, the disorder preferably is HER2-expressing cancer, *e.g.* a benign or malignant tumor characterized by overexpression of the HER2 receptor. Such cancers include, but are not limited to, breast cancer, squamous cell cancer, small-cell lung
15 cancer, non-small cell lung cancer, gastrointestinal cancer, pancreatic cancer, glioblastoma, cervical cancer, ovarian cancer, bladder cancer, hepatoma, colon cancer, colorectal cancer, endometrial carcinoma, salivary gland carcinoma, kidney cancer, liver cancer, prostate cancer, vulval cancer, thyroid cancer, hepatic carcinoma and various types of head and neck cancer.

According to the teachings herein, one may prepare a polypeptide with a variant Fc region
20 which has improved, or diminished, ADCC activity. Such molecules will find applications in the treatment of different disorders.

For example, the polypeptide variant with improved ADCC activity may be employed in the treatment of diseases or disorders where destruction or elimination of tissue or foreign micro-organisms is desired. For example, the polypeptide may be used to treat cancer; inflammatory
25 disorders; infections (*e.g.* bacterial, viral, fungal or yeast infections); and other conditions (such as goiter) where removal of tissue is desired, etc.

Where the polypeptide variant has diminished ADCC activity, such variants may be used to treat diseases or disorders where a Fc region-containing polypeptide with long half-life is desired, but the polypeptide preferably does not have undesirable effector function(s). For
30 example, the Fc region-containing polypeptide may be an anti-tissue factor (TF) antibody; anti-IgE antibody; and anti-integrin antibody (*e.g.* an anti- $\alpha 4\beta 7$ antibody). The desired mechanism of action of such Fc region-containing polypeptides may be to block ligand-receptor binding pairs. Moreover, the Fc-region containing polypeptide with diminished ADCC activity may be an agonist antibody.

The polypeptide variant is administered by any suitable means, including parenteral, subcutaneous, intraperitoneal, intrapulmonary, and intranasal, and, if desired for local immunosuppressive treatment, intralesional administration. Parenteral infusions include intramuscular, intravenous, intraarterial, intraperitoneal, or subcutaneous administration. In addition, the polypeptide variant is suitably administered by pulse infusion, particularly with declining doses of the polypeptide variant. Preferably the dosing is given by injections, most preferably intravenous or subcutaneous injections, depending in part on whether the administration is brief or chronic.

For the prevention or treatment of disease, the appropriate dosage of polypeptide variant will depend on the type of disease to be treated, the severity and course of the disease, whether the polypeptide variant is administered for preventive or therapeutic purposes, previous therapy, the patient's clinical history and response to the polypeptide variant, and the discretion of the attending physician. The polypeptide variant is suitably administered to the patient at one time or over a series of treatments.

Depending on the type and severity of the disease, about 1 $\mu\text{g/kg}$ to 15 mg/kg (e.g., 0.1-20 mg/kg) of polypeptide variant is an initial candidate dosage for administration to the patient, whether, for example, by one or more separate administrations, or by continuous infusion. A typical daily dosage might range from about 1 $\mu\text{g/kg}$ to 100 mg/kg or more, depending on the factors mentioned above. For repeated administrations over several days or longer, depending on the condition, the treatment is sustained until a desired suppression of disease symptoms occurs. However, other dosage regimens may be useful. The progress of this therapy is easily monitored by conventional techniques and assays.

The polypeptide variant composition will be formulated, dosed, and administered in a fashion consistent with good medical practice. Factors for consideration in this context include the particular disorder being treated, the particular mammal being treated, the clinical condition of the individual patient, the cause of the disorder, the site of delivery of the agent, the method of administration, the scheduling of administration, and other factors known to medical practitioners.

The "therapeutically effective amount" of the polypeptide variant to be administered will be governed by such considerations, and is the minimum amount necessary to prevent, ameliorate, or treat a disease or disorder. The polypeptide variant need not be, but is optionally formulated with one or more agents currently used to prevent or treat the disorder in question. The effective amount of such other agents depends on the amount of polypeptide variant present in the formulation, the type of disorder or treatment, and other factors discussed above. These are generally used in the same dosages and with administration routes as used hereinbefore or about from 1 to 99% of the heretofore employed dosages.

The invention will be more fully understood by reference to the following examples. They should not, however, be construed as limiting the scope of this invention. All literature and patent citations mentioned herein are expressly incorporated by reference.

EXAMPLE 1

5 Low Affinity Receptor Binding Assay

This assay determines binding of an IgG Fc region to recombinant FcγRIIA, FcγRIIB and FcγRIIA α subunits expressed as His6-glutathione S transferase (GST)-tagged fusion proteins. Since the affinity of the Fc region of IgG1 for the FcγRI is in the nanomolar range, the binding of IgG1 Fc variants can be measured by titrating monomeric IgG and measuring bound IgG with a polyclonal anti-IgG in a standard ELISA format (Example 2 below). The affinity of the other members of the FcγR family, *i.e.* FcγRIIA, FcγRIIB and FcγRIIA for IgG is however in the micromolar range and binding of monomeric IgG1 for these receptors can not be reliably measured in an ELISA format.

The following assay utilizes Fc variants of recombinant anti-IgE E27 (Figures 4A and 4B) which, when mixed with human IgE at a 1:1 molar ratio, forms a stable hexamer consisting of three anti-IgE molecules and three IgE molecules. A recombinant chimeric form of IgE (chimeric IgE) was engineered and consists of a human IgE Fc region and the Fab of an anti-VEGF antibody (Presta *et al. Cancer Research* 57:4593-4599 (1997)) which binds two VEGF molecules per mole of anti-VEGF. When recombinant human VEGF is added at a 2:1 molar ratio to chimeric IgE:E27 hexamers, the hexamers are linked into larger molecular weight complexes via the chimeric IgE Fab:VEGF interaction. The E27 component of this complex binds to the FcγRIIA, FcγRIIB and FcγRIIA α subunits with higher avidity to permit detection in an ELISA format.

MATERIALS AND METHODS

Receptor Coat: Fcγ receptor α subunits were expressed as GST fusions of His6 tagged extracellular domains (ECDs) in 293 cells resulting in an ECD-6His-GST fusion protein (Graham *et al. J. Gen. Virol.* 36:59-74 (1977) and Gorman *et al. DNA Prot. Eng. Tech.* 2:3-10 (1990)) and purified by Ni-NTA column chromatography (Qiagen, Australia) and buffer exchanged into phosphate buffered saline (PBS). Concentrations were determined by absorption at 280nm using extinction coefficients derived by amino acid composition analysis. Receptors were coated onto Nunc F96 maxisorb plates (cat no. 439454) at 100ng per well by adding 100 μl of receptor-GST fusion at 1 μg/ml in PBS and incubated for 48 hours at 4°C. Prior to assay, plates are washed 3x with 250 μl of wash buffer (PBS pH 7.4 containing 0.5% TWEEN 20™) and blocked with 250 μl of assay buffer (50mM Tris buffered saline, 0.05% TWEEN 20™, 0.5% RIA grade bovine albumin (Sigma A7888), and 2mM EDTA pH 7.4).

Immune Complex Formation: Equal molar amounts (1:1) of E27 and recombinant chimeric IgE which binds two moles recombinant human VEGF per mole of chimeric IgE are added to a 12 x 75mm polypropylene tube in PBS and mixed by rotation for 30 minutes at 25°C. E27 (anti-IgE) /chimeric IgE (IgE) hexamers are formed during this incubation. Recombinant human VEGF (165 form, MW 44,000) is added at a 2:1 molar ratio to the IgE concentration and mixed by rotation an additional 30 minutes at 25°C. VEGF- chimeric IgE binding links E27:chimeric IgE hexamers into larger molecular weight complexes which bind FcγR α subunit ECD coated plates via the Fc region of the E27 antibody.

E27:chimeric IgE:VEGF: (1:1:2 molar ratio) complexes are added to FcγR α subunit coated plates at E27 concentrations of 5 μg and 1 μg total IgG in quadruplicate in assay buffer and incubated for 120 minutes at 25°C on an orbital shaker.

Complex Detection: Plates are washed 5x with wash buffer to remove unbound complexes and IgG binding is detected by adding 100 μl horse radish peroxidase (HRP) conjugated goat anti-human IgG (γ) heavy chain specific (Boehringer Mannheim 1814249) at 1:10,000 in assay buffer and incubated for 90min at 25°C on an orbital shaker. Plates are washed 5x with wash buffer to remove unbound HRP goat anti-human IgG and bound anti-IgG is detected by adding 100 μl of substrate solution (0.4mg/ml o-phenylenedaimine dihydrochloride, Sigma P6912, 6 mM H₂O₂ in PBS) and incubating for 8 min at 25°C. Enzymatic reaction is stopped by the addition of 100 μl 4.5N H₂SO₄ and colorimetric product is measured at 490 nm on a 96 well plate densitometer (Molecular Devices). Binding of E27 variant complexes is expressed as a percent of the wild type E27 containing complex.

EXAMPLE 2

Identification of Unique C1q Binding Sites in a Human IgG Antibody

In the present study, mutations were identified in the CH2 domain of a human IgG1 antibody, "C2B8" (Reff *et al.*, *Blood* 83:435 (1994)), that ablated binding of the antibody to C1q but did not alter the conformation of the antibody nor affect binding to each of the FcγRs. By alanine scanning mutagenesis, five variants in human IgG1 were identified, D270K, D270V, K322A P329A, and P331, that were non-lytic and had decreased binding to C1q. The data suggested that the core C1q binding sites in human IgG1 is different from that of murine IgG2b. In addition, K322A, P329A and P331A were found to bind normally to the CD20 antigen, and to four Fc receptors, FcγRI, FcγRII, FcγRIII and FcRn.

MATERIALS AND METHODS

Construction of C2B8 Variants: The chimeric light and heavy chains of anti-CD20 antibody C2B8 (Reff *et al.*, *Blood* 83:435 (1994)) subcloned separately into previously described PRK vectors (Gorman *et al.*, *DNA Protein Eng. Tech.* 2:3 (1990)) were used. By site directed

mutagenesis (Kunkel *et al.*, *Proc. Natl. Acad. Sci. USA* 82:488 (1987)), alanine scan variants of the Fc region in the heavy chain were constructed. The heavy and light chain plasmids were co-transfected into an adenovirus transformed human embryonic kidney cell line as previously described (Werther *et al.*, *J. Immunol.* 157:4986 (1996)). The media was changed to serum-free
5 24 hours after transfection and the secreted antibody was harvested after five days. The antibodies were purified using Protein A-SEPHAROSE CL-4B™ (Pharmacia), buffer exchanged and concentrated to 0.5 ml with PBS using a Centricon-30 (Amicon), and stored at 4°C. The concentration of the antibody was determined using total Ig-binding ELISA.

C1q Binding ELISA: Costar 96 well plates were coated overnight at 4°C with the indicated
10 concentrations of C2B8 in coating buffer (0.05 M sodium carbonate buffer), pH 9. The plates were then washed 3x with PBS/ 0.05% TWEEN 20™, pH 7.4 and blocked with 200µl of ELISA diluent without thimerosal (0.1M NaPO₄ / 0.1M NaCl / 0.1% gelatin / 0.05% TWEEN 20™/ 0.05% ProClin300) for 1hr at room temperature. The plate was washed 3x with wash buffer, an aliquot of 100µl of 2 µg/ml C1q (Quidel, San Diego, CA) was added to each well and incubated for 2 hrs
15 at room temperature. The plate was then washed 6x with wash buffer. 100 µl of a 1:1000 dilution of sheep anti-complement C1q peroxidase conjugated antibody (Bioscience) was added to each well and incubated for 1 hour at room temperature. The plate was again washed 6x with wash buffer and 100 µl of substrate buffer (PBS/0.012% H₂O₂) containing OPD (O-phenylenediamine dihydrochloride (Sigma)) was added to each well. The oxidation reaction, observed by the
20 appearance of a yellow color, was allowed to proceed for 30 minutes and stopped by the addition of 100 µl of 4.5 N H₂SO₄. The absorbance was then read at (492-405) nm using a microplate reader (SPECTRA MAX 250™, Molecular Devices Corp.). The appropriate controls were run in parallel (*i.e.* the ELISA was performed without C1q for each concentration of C2B8 used and also the ELISA was performed without C2B8). For each variant, C1q binding was measured by
25 plotting the absorbance (492-405) nm versus concentration of C2B8 in µg/ml using a 4-parameter curve fitting program (KALEIDAGRAPH™) and comparing EC₅₀ values.

Complement Dependent Cytotoxicity (CDC) Assay: This assay was performed essentially as previously described (Gazzano-Santoro *et al.*, *J. Immunol. Methods* 202:163 (1996)). Various concentrations of C2B8 (0.08-20 µg/ml) were diluted with RHB buffer (RPMI
30 1640/20mM HEPES (pH 7.2)/2mM Glutamine/0.1% BSA/100 µg/ml Gentamicin). Human complement (Quidel) was diluted 1:3 in RHB buffer and WIL2-S cells (available from the ATCC, Manassas, VA) which express the CD20 antigen were diluted to a density of 1 x 10⁶ cells /ml with RHB buffer. Mixtures of 150 µl containing equal volumes of C2B8, diluted human complement and WIL2-S cells were added to a flat bottom tissue culture 96 well plate and allowed to incubate
35 for 2 hrs at 37°C and 5% CO₂ to facilitate complement mediated cell lysis. 50 µl of alamar blue

(Accumed International) was then added to each well and incubated overnight at 37°C. The absorbance was measured using a 96-well fluorometer with excitation at 530 nm and emission at 590 nm. As described by Gazzano-Santoro *et al.*, the results are expressed in relative fluorescence units (RFU). The sample concentrations were computed from a C2B8 standard curve and the percent activity as compared to wild type C2B8 is reported for each variant.

CD20 Binding Potency of the C2B8 Variants: The binding of C2B8 and variants to the CD20 antigen were assessed by a method previously described (Reff *et al.*, (1994), *supra*; reviewed in Gazzano-Santoro *et al.*, (1996), *supra*). WIL2-S cells were grown for 3-4 days to a cell density of 1×10^6 cells/ml. The cells were washed and spun twice in FACS buffer (PBS/0.1% BSA/0.02% NaN_3) and resuspended to a cell density of 5×10^6 cells/ml. 200 μl of cells (5×10^6 cells/ml) and 20 μl of diluted C2B8 samples were added to a 5 ml tube and incubated at room temperature for 30 minutes with agitation. The mixture was then washed with 2 ml of cold FACS buffer, spun down and resuspended in 200 μl of cold FACS buffer. To the suspension, 10 μl of goat anti-human IgG-FITC (American Qualex Labs.) was added and the mixture was incubated in the dark at room temperature for 30 minutes with agitation. After incubation, the mixture was washed with 2 ml of FACS buffer, spun down and resuspended in 1 ml of cold fixative buffer (1% formaldehyde in PBS). The samples were analyzed by flow cytometry and the results expressed as relative fluorescence units (RFU) were plotted against antibody concentrations using a 4-parameter curve fitting program (KALEIDAGRAPH™). The EC_{50} values are reported as a percentage of that of the C2B8 reference material.

Fc γ R Binding ELISAs: Fc γ RI α subunit-GST fusion was coated onto Nunc F96 maxisorb plates (cat no. 439454) by adding 100 μl of receptor-GST fusion at 1 $\mu\text{g}/\text{ml}$ in PBS and incubated for 48 hours at 4°C. Prior to assay, plates are washed 3x with 250 μl of wash buffer (PBS pH 7.4 containing 0.5% TWEEN 20™) and blocked with 250 μl of assay buffer (50mM Tris buffered saline, 0.05% TWEEN 20™, 0.5% RIA grade bovine albumin (Sigma A7888), and 2mM EDTA pH 7.4). Samples diluted to 10 $\mu\text{g}/\text{ml}$ in 1 ml of assay buffer are added to Fc γ RI α subunit coated plates and incubated for 120 minutes at 25°C on an orbital shaker. Plates are washed 5x with wash buffer to remove unbound complexes and IgG binding is detected by adding 100 μl horse radish peroxidase (HRP) conjugated goat anti-human IgG (γ) heavy chain specific (Boehringer Mannheim 1814249) at 1:10,000 in assay buffer and incubated for 90min at 25°C on an orbital shaker. Plates are washed 5x with wash buffer to remove unbound HRP goat anti-human IgG and bound anti-IgG is detected by adding 100 μl of substrate solution (0.4mg/ml o-phenylenediamine dihydrochloride, Sigma P6912, 6 mM H_2O_2 in PBS) and incubating for 8 min at 25°C. Enzymatic reaction is stopped by the addition of 100 μl 4.5N H_2SO_4 and colorimetric

product is measured at 490 nm on a 96 well plate densitometer (Molecular Devices). Binding of variant is expressed as a percent of the wild type molecule.

FcγRII and III binding ELISAs were performed as described in Example 1 above.

For measuring FcRn binding activity of IgG variants, ELISA plates were coated with 2
 5 μg/ml streptavidin (Zymed, South San Francisco) in 50 mM carbonate buffer, pH 9.6, at 4°C
 overnight and blocked with PBS-0.5% BSA, pH 7.2 at room temperature for one hour. Biotinylated
 FcRn (prepared using biotin-X-NHS from Research Organics, Cleveland, OH and used at 1-2
 μg/ml) in PBS-0.5% BSA, 0.05% polysorbate 20, pH 7.2, was added to the plate and incubated
 for one hour. Two fold serial dilutions of IgG standard (1.6-100 ng/ml) or variants in PBS-0.5%
 10 BSA, 0.05% polysorbate 20, pH 6.0, were added to the plate and incubated for two hours. Bound
 IgG was detected using peroxidase labeled goat F(ab')₂ anti-human IgG F(ab')₂ in the above pH
 6.0 buffer (Jackson ImmunoResearch, West Grove, PA) followed by 3,3',5,5'-tetramethyl
 benzidine (Kirgaard & Perry Laboratories) as the substrate. Plates were washed between steps
 with PBS-0.05% polysorbate 20 at either pH 7.2 or 6.0. Absorbance was read at 450 nm on a
 15 Vmax plate reader (Molecular Devices, Menlo Park, CA). Titration curves were fit with a four-
 parameter nonlinear regression curve-fitting program (KaleidaGraph, Synergy software, Reading,
 PA). Concentrations of IgG variants corresponding to the mid-point absorbance of the titration
 curve of the standard were calculated and then divided by the concentration of the standard
 corresponding to the mid-point absorbance of the standard titration curve.

RESULTS AND DISCUSSION

By alanine scanning mutagenesis, several single point mutations were constructed in the
 CH2 domain of C2B8 beginning with E318A, K320A and K322A. All the variants constructed
 bound normally to the CD20 antigen (Table 3).

Table 3

	wt	E318A	K320A	K322A	P329A	P331A
FcRn	+	+	+	+		
CD20	+	+	+	+	+	+
FcγRI	+	+	+	+	+	+
FcγRII	+	+	+	+	+	+
FcγRIII	+	+	+	+	+	+
*C1q	+++	++	+++	-	-	-
CDC	+	+	+	-	-	-

25 (+) indicates binding and (-) signifies binding abolished

*With respect to C1q binding, each + sign is equivalent to approximately 33% binding.

Where binding of human complement to an antibody with a human Fc was analyzed, the ability of E318A and K320A to activate complement was essentially identical to that of wild type C2B8 (Table 3). When compared to wild type C2B8, there appears to be little difference in the binding of E318A and K320A to C1q. There is only a 10% decrease in the binding of K320A and about a 30% decrease in the binding of E318A to C1q (Fig. 2). The results indicate that the effect of the E318A and the K320A substitution on complement activation and C1q binding is minimal. Also, the human IgG1 of C2B8 was substituted for human IgG2 and used as a negative control in the C1q binding studies. The IgG2 variant appears to have a much lower affinity for C1q than the E318A and K320A variants (Fig. 2). Thus, the results demonstrate that E318 and K320 do not constitute the core C1q binding sites for human IgG1. Conversely, the K322A substitution had a significant effect on both complement activity and C1q binding. The K322A variant did not have CDC activity when tested in the above CDC assay and was more than a 100 fold lower than wild type C2B8 in binding to C1q (Fig. 2). In the human system, K322 is the only residue of the proposed core C1q binding sites that appeared to have a significant effect on complement activation and C1q binding.

Since the Duncan and Winter study was performed using mouse IgG2b and the above results reveal that K320 and E318 in human IgG1 are not involved in C1q binding, and without being bound to any one theory, the above data suggest that the C1q binding region in murine IgGs is different from that of the human. To investigate this further and also to identify additional variants that do not bind to C1q and hence do not activate complement, several more point mutations in the vicinity of K322 were constructed as assessed from the three dimensional structure of the C2B8 Fc. Variants constructed, K274A, N276A, Y278A, S324A, P329A, P331A, K334A, and T335A, were assessed for their ability to bind C1q and also to activate complement. Many of these substitutions had little or no effect on C1q binding or complement activation. In the above assays, the P329A and the P331A variants did not activate complement and had decreased binding to C1q. The P331A variant did not activate complement and was 60 fold lower in binding to C1q (Fig. 3) when compared to wild type C2B8 (Fig. 2). The concentration range of the antibody variants used in Fig. 3 is expanded to 100µg/ml in order to observe saturation of C1q binding to the P331A variant. The mutation P329A results in an antibody that does not activate complement and is more than a 100 fold lower in binding to C1q (Fig. 3) when compared to wild type C2B8 (Fig. 2).

Variants that did not bind to C1q and hence did not activate complement were examined for their ability to bind to the Fc receptors: FcγRI, FcγRIIA, FcγRIIB, FcγRIIIA and FcRn. This particular study was performed using a humanized anti-IgE antibody, an IgG1 antibody with these mutations (see Example 1 above). The results revealed the variants, K322A and P329A, bind

to all the Fc receptors to the same extent as the wild type protein (Table 4). However, there was a slight decrease in the binding of P331A to FcγRIIB.

In conclusion, two amino acid substitutions in the COOH terminal region of the CH2 domain of human IgG1, K322A and P329A were identified that result in more than 100 fold decrease in C1q binding and do not activate the CDC pathway. These two variants, K322A and P329A, bind to all Fc receptors with the same affinity as the wild type antibody. Based on the results, summarized in Table 4, and without being bound to any one theory, it is proposed that the C1q binding epicenter of human IgG1 is centered around K322, P329 and P331 and is different from the murine IgG2b epicenter which constitutes E318, K320 and K322.

Table 4

	wt	E318A	K320A	K322A	P329A	P331A
CD20	100	89	102	86	112	103
^a FcγRI	100	93	102	90	104	74
^a FcγRIIA	100	113	94	109	111	86
^a FcγRIIB	100	106	83	101	96	58
^a FcγRIII	100	104	72	90	85	73
CDC	100	108	108	none	none	none

^aFor binding to the FcγRs the variants were made in the E27 background (anti-IgE). The results are presented as a percentage of the wild type.

A further residue involved in binding human C1q was identified using the methods described in the present example. The residue D270 was replaced with lysine and valine to generate variants D270K and D270V, respectively. These variants both showed decreased binding to human C1q (Fig. 6) and were non-lytic (Fig. 7). The two variants bound the CD20 antigen normally and recruited ADCC.

EXAMPLE 3

Variants with Improved C1q Binding

The following study shows that substitution of residues at positions K326, A327, E333 and K334 resulted in variants with at least about a 30% increase in binding to C1q when compared to the wild type antibody. This indicated K326, A327, E333 and K334 are potential sites for improving the efficacy of antibodies by way of the CDC pathway. The aim of this study was to improve CDC activity of an antibody by increasing binding to C1q. By site directed mutagenesis at K326 and E333, several variants with increased binding to C1q were constructed. The residues in order of increased binding at K326 are K<V<E<A<G<D<M<W, and the residues in order of increased binding at E333 are E<Q<D<V<G<A<S. Four variants, K326M, K326D, K326E

and E333S were constructed with at least a two-fold increase in binding to C1q when compared to wild type. Variant K326W displayed about a five-fold increase in binding to C1q.

5 Variants of the wild type C2B8 antibody were prepared as described above in Example 2. A further control antibody, wild type C2B8 produced in Chinese hamster ovary (CHO) cells essentially as described in US Patent 5,736,137, was included in a C1q binding ELISA to confirm that wt C2B8 produced in the 293 kidney cell line had the same C1q binding activity as the CHO-produced antibody (see "CHO-wt-C2B8" in Fig. 8). The C1q binding ELISA, CDC assay, and CD20 binding potency assay in this example were performed as described in Example 2 above.

10 As shown in Fig. 8, alanine substitution at K326 and E333 in C2B8 resulted in variants with about a 30% increase in binding to C1q.

Several other single point variants at K326 and E333 were constructed and assessed for their ability to bind C1q and activate complement. All the variants constructed bound normally to the CD20 antigen.

15 With respect to K326, the other single point variants constructed were K326A, K326D, K326E, K326G, K326V, K326M and K326W. As shown in Fig. 9, these variants all bound to C1q with a better affinity than the wild type antibody. K326W, K326M, K326D and K326E showed at least a two-fold increase in binding to C1q (Table 5). Among the K326 variants, K326W had the best affinity for C1q.

Table 5

Variant	EC ₅₀ value
Wild type	1.53
K326V	1.30
K326A	1.03
K326E	1.08
K326G	0.95
K326D	0.76
K326M	0.67
K326W	0.47
E333S	0.81
E333A	0.98
E333G	1.14
E333V	1.18
E333D	1.22
E333Q	1.52
K334A	1.07

Substitutions with hydrophobic as well as charged residues resulted in variants with increased binding to C1q. Even substitution with glycine which is known to impart flexibility to a chain and is well conserved in nature, resulted in a variant with higher affinity for C1q when compared to the wild type. It would appear that any amino acid substitution at this site would result in a variant with higher affinity for C1q. As assessed from the three-dimensional structure, K326 and E333 are in the vicinity of the core C1q binding sites (Fig. 10).

In addition to alanine, E333 was also substituted with other amino acid residues. These variants, E333S, E333G, E333V, E333D, and E333Q, all had increased binding to C1q when compared to the wild type (Fig. 11). As shown in Table 5, the order of binding affinity for C1q was as follows: E333S>E333A>E333G>E333V>E333D>E333Q. Substitutions with amino acid residues with small side chain volumes, *i.e.* serine, alanine and glycine, resulted in variants with higher affinity for C1q in comparison to the other variants, E333V, E333D and E333Q, with larger side chain volumes. The variant E333S had the highest affinity for C1q, showing a two-fold increase in binding when compared to the wild type. Without being bound to any one theory, this indicates the effect on C1q binding at 333 may also be due in part to the polarity of the residue.

Double variants were also generated. As shown in Figs. 12 and 13, double variants K326M-E333S and K326A-E333A were at least three-fold better at binding human C1q than wild type C2B8 (Fig. 12) and at least two-fold better at mediating CDC compared to wild type C2B8 (Fig. 13). Additivity indicates these are independently acting variants.

As shown in Fig. 14, a further variant with improved C1q binding (50% increase) was made by changing A327 in a human IgG1 constant region to glycine. Conversely, in a human IgG2 constant region, changing G327 to alanine reduced C1q binding of the IgG2 antibody.

EXAMPLE 4

Identification of FcR Binding Sites in Human IgG Antibodies

In the present study, the effect of mutating various Fc region residues of an IgG1 antibody with respect to binding FcγRI, FcγRIIA, FcγRIIB and FcγRIIIA as well as FcRn was evaluated. Antibody variants with improved as well as diminished FcR binding were identified.

MATERIALS AND METHODS

Construction of IgG1 Variants: Recombinant anti-IgE E27 having the light chain and heavy chain sequences in Figs. 4A and 4B, respectively, was used as the parent antibody in the following experiments. This antibody binds the antigen IgE and has a non-A allotype IgG1 Fc region. By site directed mutagenesis (Kunkel *et al.*, *Proc. Natl. Acad. Sci. USA* 82:488 (1987)), variants of the Fc region in the heavy chain of the above parent antibody were constructed. The heavy and light chain plasmids were co-transfected into an adenovirus transformed human

embryonic kidney cell line as previously described (Werther *et al.*, *J. Immunol.* 157:4986 (1996)). The media was changed to serum-free 24 hours after transfection and the secreted antibody was harvested after five days. The antibodies were purified by Protein G SEPHAROSE® (Pharmacia), buffer exchanged and concentrated to 0.5 ml with PBS using a Centricon-30 (Amicon), and stored at 4°C. Concentration was determined by adsorption at 280 nm using extinction coefficients derived by amino acid composition analysis.

High Affinity Fc γ RIA Binding ELISA: Fc γ RIA was expressed as a GST fusion of His6 tagged extracellular domain in 293 cells and purified by Ni-NTA column chromatography.

To purify Fc γ RIA, supernatant from transfected 293 cells was removed after three days. Protease inhibitors were added; 50 μ L Aprotinin (Sigma)/ 50 mL supernatant, and PMSF (1mM). Supernatants were concentrated to 10 mL in a stirred cell (Amicon), and dialyzed overnight at 4°C against 1 liter column buffer (50 mM Tris pH 8.0, 20mM Imidazole, 300 mM NaCl). Additional dialysis was done the following morning against fresh column buffer for 4 hours at 4°C. The solution was loaded on to a 1mL Ni⁺⁺ column (NTA super flow resin, Qiagen) previously equilibrated with 10 mL column buffer. Columns were washed with 10 mL column buffer, and protein was eluted with 2.5 mL elution buffer (50mM Tris pH 8.0, 250mM Imidazole, 300mM NaCl). Protein was concentrated to 0.5 mL and buffer exchanged into PBS. Concentrations were determined by adsorption at 280 nm using an extinction coefficient derived by amino acid composition analysis.

Purified receptors were coated onto Nunc F96 maxisorb plates (cat no. 439545) at approximately 150 ng per well by adding 100 μ L of receptor at 1.5 μ g/mL in PBS and incubated for 24 hours at 4°C. Prior to assay, plates were washed 3x with 250 μ L of wash buffer (phosphate buffered saline pH 7.4 containing 0.5% TWEEN 20®) and blocked with 250 μ L of assay buffer (50 mM tris buffered saline, 0.05% TWEEN 20®, 0.5% RIA grade bovine albumin (Sigma A7888), and 2mM EDTA pH 7.4).

100 μ L of E27 was added to the first four wells of the Fc γ RIA subunit coated plated at a concentration of 10 μ g/mL. 80 μ L of assay buffer was added to the next four well followed by 20 μ L of the 10 μ g/mL E27 IgG to give a final concentration of 2 μ g/mL. Plates were incubated at 25°C for 2 hours on an orbital shaker.

For detection, plates were washed 5x with wash buffer to remove unbound antibody. IgG binding to GST-Fc γ RIA was detected by adding 100 μ L horse radish peroxidase (HRP) conjugated protein G (BIORAD) at 1:5000. HRP conjugates were incubated for 1.5 hours at 25°C on an orbital shaker. Plates were washed x5 with wash buffer to remove unbound HRP conjugate. Binding was detected by adding 100 μ L of substrate solution (0.4mg/mL o-

phenylenedaimine dihydrochloride, Sigma P6912, 6 mM H₂O₂ in PBS) and incubating for 10 minutes at 25°C. Enzymatic reaction was stopped by the addition of 100 µL of 4.5 N H₂SO₄ and colorimetric product was measured at 490 nm on a 96 well plate densitometer (Molecular Devices).

5 Binding of E27 variants at IgG concentration of 2 µg/mL was expressed as a ratio of wild type E27.

FcγRIA THP-1 Assay. 100 µL of E27 was added to the first three wells of a serocluster plate (Costar) at a concentration of 20 µg/mL in assay buffer (1x PBS, 0.1% BSA, 0.01% NaN₃). 92.5 µL of assay buffer was added to the next three wells followed by 7.5 µL of the 20µg/mL E27 IgG to give a final concentration of 1.5 µg/mL. To each well, 100 µL of THP-1 cells were added at a concentration of 5 million cells/mL in FACS assay buffer. The plate is incubated on ice for 10 30 minutes

For detection, cells were washed 2x with assay buffer to remove unbound antibody. IgG binding FcγRIA was detected by adding 100 µL FITC conjugated F(ab')₂ fragment of goat anti-human IgG heavy chain specific. (Jackson ImmunoResearch) at 1:200. FITC conjugates were 15 incubated with cells for 30 minutes on ice. Cells were washed x3 with assay buffer to remove unbound FITC conjugate. Cells were stained with P.I. (SIGMA) at 2.5 µg/mL and analyzed by flow cytometry.

Binding of E27 variants at IgG concentration of 1.5 µg/mL was expressed as a ratio of wild 20 type E27.

Data from the plate assay (FcγRIA ELISA) and cell-based assay (FcγRIA THP-1 assay) was averaged to arrive at an FcγRIA-binding activity.

Low Affinity FcγR Binding ELISAs: FcγRIIA, FcγRIIB and FcγRIIIA binding ELISAs were performed as described in Example 1 above, with detection of the stable hexamer (consisting of 25 three anti-IgE molecules and three IgE molecules).

FcRn Binding ELISA: For measuring FcRn binding activity of IgG variants, ELISA plates were coated with 2 µg/ml streptavidin (Zymed, South San Francisco) in 50 mM carbonate buffer, pH 9.6, at 4°C overnight and blocked with PBS-0.5% BSA, pH 7.2 at room temperature for one hour. Biotinylated FcRn (prepared using biotin-X-NHS from Research Organics, Cleveland, OH and used at 1-2 µg/ml) in PBS-0.5% BSA, 0.05% polysorbate 20, pH 7.2, was added to the plate 30 and incubated for one hour. Two fold serial dilutions of IgG standard (1.6-100 ng/ml) or variants in PBS-0.5% BSA, 0.05% polysorbate 20, pH 6.0, were added to the plate and incubated for two hours. Bound IgG was detected using peroxidase labeled goat F(ab')₂ anti-human IgG F(ab')₂ in the above pH 6.0 buffer (Jackson ImmunoResearch, West Grove, PA) followed by 3,3',5,5'- 35 tetramethyl benzidine (Kirgaard & Perry Laboratories) as the substrate. Plates were washed

between steps with PBS-0.05% TWEEN 20® at either pH 7.2 or 6.0. Absorbance was read at 450 nm on a Vmax plate reader (Molecular Devices, Menlo Park, CA). Titration curves were fit with a four-parameter nonlinear regression curve-fitting program (KaleidaGraph, Synergy software, Reading, PA). Concentrations of IgG variants corresponding to the mid-point absorbance of the titration curve of the standard were calculated and then divided by the concentration of the standard corresponding to the mid-point absorbance of the standard titration curve.

In Vitro ADCC Assay: To prepare chromium 51-labeled target cells, tumor cell lines were grown in tissue culture plates and harvested using sterile 10 mM EDTA in PBS. SK-BR-3 cells, a 3+ HER2-overexpressing human breast cancer cell line, were used as targets in all assays. The detached cells were washed twice with cell culture medium. Cells (5×10^6) were labeled with 200 μ Ci of chromium 51 (New England Nuclear/DuPont) at 37°C for one hour with occasional mixing. Labeled cells were washed three times with cell culture medium, then were resuspended to a concentration of 1×10^5 cells/mL. Cells were used either without opsonization, or were opsonized prior to the assay by incubation with rhuMAb HER2 wildtype (HERCEPTIN®) or seven Fc mutants (G14, G18, G17, G36, G30, G31 and G34) at 100 ng/mL and 1.25 ng/mL in PBMC assay or 20 ng/mL and 1 ng/mL in NK assay.

Peripheral blood mononuclear cells were prepared by collecting blood on heparin from normal healthy donors and dilution with an equal volume of phosphate buffered saline (PBS). The blood was then layered over LYMPHOCYTE SEPARATION MEDIUM® (LSM: Organon Teknika) and centrifuged according to the manufacturer's instructions. Mononuclear cells were collected from the LSM-plasma interface and were washed three times with PBS. Effector cells were suspended in cell culture medium to a final concentration of 1×10^7 cells/mL.

After purification through LSM, natural killer (NK) cells were isolated from PBMCs by negative selection using an NK cell isolation kit and a magnetic column (Miltényi Biotech) according to the manufacturer's instructions. Isolated NK cells were collected, washed and resuspended in cell culture medium to a concentration of 2×10^6 cells/mL. The identity of the NK cells was confirmed by flow cytometric analysis.

Varying effector:target ratios were prepared by serially diluting the effector (either PBMC or NK) cells two-fold along the rows of a microtiter plate (100 μ L final volume) in cell culture medium. The concentration of effector cells ranged from 1.0×10^7 /mL to 2.0×10^4 /mL for PBMC and from 2.0×10^6 /mL to 3.9×10^3 /mL for NK. After titration of effector cells, 100 μ L of chromium 51-labeled target cells (opsonized or nonopsonized) at 1×10^5 cells/mL were added to each well of the plate. This resulted in an initial effector:target ratio of 100:1 for PBMC and 20:1 for NK cells. All assays were run in duplicate, and each plate contained controls for both spontaneous lysis (no

effector cells) and total lysis (target cells plus 100 μ L) 1% sodium dodecyl sulfate, 1 N sodium hydroxide). The plates were incubated at 37°C for 18 hours, after which the cell culture supernatants were harvested using a supernatant collection system (Skatron Instrument, Inc.) and counted in a Minaxi auto-gamma.5000 series gamma counter (Packard) for one minute. Results
5 were then expressed as percent cytotoxicity using the formula:

$$\% \text{ Cytotoxicity} = (\text{sample cpm} - \text{spontaneous lysis}) / (\text{total lysis} - \text{spontaneous lysis}) \times 100$$

Four-parameter curve-fitting was then used to evaluate the data (KaleidaGraph 3.0.5).

RESULTS

A variety of antibody variants were generated which had FcR binding activity that differed
10 from the parent antibody. The FcR binding data for the variants generated is shown in Tables 6 and 7 below. An additional variant, T307Q, also displayed improved FcRn binding compared to E27 parent antibody.

TABLE 6
CH2 DOMAIN VARIANTS

IG2	Res#EU (Kabat)	FcRn mean	sd	n	FcγRI mean	sd	n	FcγRIIA mean	sd	FcγRIIB mean	sd	FcγRIIa mean	sd
REDUCED BINDING TO ALL FcγR													
1	233-236	0.54	(0.20)	3	0.12	(0.06)	6	0.08	(0.01)	0.12	(0.01)	0.04	(0.02)
	ELLG > PVA-												
2	P238A(251)	1.49	(0.17)	3	0.60	(0.05)	5	0.38	(0.14)	0.36	(0.15)	0.07	(0.05)
14	D265A(278)	1.23	(0.14)	4	0.14	(0.04)	6	0.07	(0.01)	0.13	(0.05)	0.09	(0.06)
17	E269A(282)	1.05			0.52	(0.03)	6	0.65	(0.18)	0.75	(0.29)	0.45	(0.13)
18	D270A(283)	1.05			0.76	(0.12)	6	0.06	(0.01)	0.11	(0.05)	0.14	(0.04)
58	N297A(314)	0.80	(0.18)	8	0.15	(0.06)	7	0.05	(0.00)	0.10	(0.02)	0.03	(0.01)
52	A327Q(346)	0.97			0.63	(0.15)	7	0.13	(0.03)	0.14	(0.03)	0.06	(0.01)
64	P329A(348)	0.80			0.48	(0.10)	6	0.08	(0.02)	0.12	(0.08)	0.21	(0.03)
REDUCED BINDING TO FcγRII & FcγRIII													
3	S239A(252)	1.06			0.81	(0.09)	7	0.73	(0.25)	0.76	(0.36)	0.26	(0.08)
33	E294A(311)	0.75			0.90	(0.08)	4	0.87	(0.19)	0.63	(0.17)	0.66	(0.14)
34	Q295A(312)	0.79			1.00	(0.11)	4	0.62	(0.20)	0.50	(0.24)	0.25	(0.09)
39	V303A(322)	1.26	(0.21)	3	0.91	(0.11)	5	0.86	(0.10)	0.65	(0.17)	0.33	(0.09)
IMPROVED BINDING TO FcγRII & FcγRIII													
11	T256A(269)	1.91	(0.43)	6	1.14	(0.14)	4	1.41	(0.27)	2.06	(0.66)	1.32	(0.18)
30	K290A(307)	0.79	(0.14)	3	1.01	(0.08)	4	1.29	(0.21)	1.40	(0.18)	1.28	(0.21)

44	D312A(331)	1.50 (0.06)	4	1.01 (0.12)	5	1.20 (0.24)	1.19 (0.07)	1.23 (0.14)	n=3
51	K326A(345)	1.03		1.04 (0.05)	4	1.26 (0.21)	1.49 (0.27)	1.22 (0.28)	n=5
197	A330(349)K					1.28	1.25	1.28	n=1
273	A339(359)T			1.23		1.11	1.23	1.42	n=1
=====									
EFFECT FcYRII									
10	R255A(268)	0.59 (0.19)	4	1.26 (0.26)	8	1.30 (0.20)	1.59 (0.42)	0.98 (0.18)	n=5
12	E258A(271)	1.18		1.18 (0.13)	4	1.33 (0.22)	1.65 (0.38)	1.12 (0.12)	n=5
15	S267A(280)	1.08		1.20 (0.14)	4	1.64 (0.18)	2.06 (0.35)	1.14 (0.25)	n=7
16	H268A(281)	1.02 (0.22)	3	1.05 (0.11)	4	1.22 (0.14)	1.45 (0.23)	0.52 (0.09)	n=12
19	E272A(285)	1.34 (0.24)	4	1.04 (0.06)	4	1.24 (0.11)	1.58 (0.19)	0.74 (0.12)	n=4
21	N276A(289)	1.15 (0.21)	3	1.05 (0.14)	4	1.29 (0.20)	1.34 (0.40)	0.95 (0.04)	n=4
23	D280A(295)	0.82		0.97 (0.06)	4	1.34 (0.14)	1.60 (0.31)	1.09 (0.20)	n=10
25	E283A(300)	0.71		0.97 (0.03)	4	1.24 (0.23)	1.20 (0.17)	1.01 (0.14)	n=5
26	H285A(302)	0.85		0.96 (0.07)	4	1.26 (0.12)	1.23 (0.15)	0.87 (0.04)	n=4
27	N286A(303)	1.24 (0.04)	2	0.94 (0.20)	13	1.28 (0.23)	1.39 (0.14)	1.03 (0.08)	n=5
31	R292A(309)	0.81 (0.18)	4	0.93 (0.02)	4	0.27 (0.14)	0.18 (0.07)	0.90 (0.18)	n=9
36	S298A(317)	0.80		1.10 (0.04)	3	0.40 (0.08)	0.21 (0.11)	1.30 (0.18)	n=12
38	R301A(320)	0.86		1.06 (0.10)	4	1.12 (0.12)	1.26 (0.14)	0.21 (0.08)	n=6
38B	R301M(320)	0.88		1.06 (0.12)	4	1.29 (0.17)	1.56 (0.12)	0.48 (0.21)	n=4
40	V305A(324)	1.46 (0.48)	6	1.04 (0.19)	10	1.12 (0.12)	1.23 (0.22)	0.84 (0.15)	n=4
41	T307A(326)	1.81 (0.32)	6	0.99 (0.14)	4	1.19 (0.37)	1.35 (0.33)	1.12 (0.18)	n=12

42	L309A(328)	0.63	(0.18)	4	0.93	(0.18)	6	1.13	(0.08)	1.26	(0.12)	1.07	(0.20)	n=3
45	N315A(334)	0.76	(0.14)	3	1.27	(0.36)	6	1.15	(0.06)	1.30	(0.17)	1.07	(0.21)	n=5
48	K320A(339)	1.10			0.98	(0.09)	5	1.12	(0.11)	1.22	(0.05)	0.87	(0.17)	n=4
49	K322A(341)	0.98			0.94	(0.05)	6	1.15	(0.11)	1.27	(0.24)	0.61	(0.14)	n=5
50	S324A(343)	1.08			0.95	(0.05)	4	0.82	(0.22)	0.70	(0.12)	1.12	(0.17)	n=4
65	P331A(350)	0.85			1.30	(0.34)	8	1.29	(0.14)	1.47	(0.28)	1.03	(0.19)	n=3
54	E333A(352)	1.03	(0.01)	2	0.98	(0.15)	5	0.92	(0.12)	0.76	(0.11)	1.27	(0.17)	n=10
56	T335A(354)	0.98			1.00	(0.05)	4	0.79	(0.22)	0.65	(0.26)	0.92	(0.54)	n=3
57	S337A(356)	1.03			1.17	(0.23)	3	1.22	(0.30)	1.26	(0.06)	0.94	(0.18)	n=4
=====														
EFFECT FcYRIII														
5	K248A(261)	0.87			0.95	(0.05)	5	1.06	(0.12)	1.01	(0.12)	0.71	(0.05)	n=4
6	D249A(262)	0.93			1.04	(0.10)	4	1.02	(0.12)	0.94	(0.02)	0.66	(0.07)	n=5
7	M252A(265)	0.64	(0.13)	4	0.99	(0.10)	5	1.01	(0.18)	1.15	(0.22)	0.65	(0.17)	n=6
9	S254A(267)	< 0.10			0.96	(0.08)	4	0.97	(0.24)	1.15	(0.38)	0.73	(0.14)	n=3
16	H268A(281)	1.02	(0.22)	3	1.05	(0.11)	4	1.22	(0.14)	1.45	(0.23)	0.52	(0.09)	n=12
19	E272A(285)	1.34	(0.24)	4	1.04	(0.06)	4	1.24	(0.11)	1.58	(0.19)	0.74	(0.12)	n=4
22	Y278A(291)	0.90			0.96	(0.02)	4	1.11	(0.08)	1.10	(0.16)	0.67	(0.11)	n=4
29	T289A(306)	0.86			0.93	(0.03)	4	0.96	(0.33)	0.83	(0.22)	0.62	(0.19)	n=7
32	E293A(310)	0.85			1.11	(0.07)	4	1.08	(0.19)	1.07	(0.20)	0.31	(0.13)	n=6
35	Y296F(313)	0.79			1.07	(0.12)	4	0.97	(0.26)	0.84	(0.18)	0.52	(0.09)	n=5
36	S298A(317)	0.80			1.10	(0.04)	3	0.40	(0.08)	0.21	(0.11)	1.30	(0.18)	n=12

38	R301A(320)	0.86	1.06 (0.10)	4	1.12 (0.12)	1.26 (0.14)	0.21 (0.08)	n=6
38B	R301M(320)	0.88	1.06 (0.12)	4	1.29 (0.17)	1.56 (0.12)	0.48 (0.21)	n=4
49	K322A(341)	0.98	0.94 (0.05)	6	1.15 (0.11)	1.27 (0.24)	0.61 (0.14)	n=5
54	E333A(352)	1.03 (0.01)	2 0.98 (0.15)	5	0.92 (0.12)	0.76 (0.11)	1.27 (0.17)	n=10
55	K334A(353)	1.05 (0.03)	2 1.10 (0.06)	4	1.01 (0.15)	0.90 (0.12)	1.39 (0.19)	n=17
NO EFFECT ON FcyR								
4	K246A(259)	1.03	0.94 (0.06)	4	1.02 (0.10)	0.92 (0.15)	1.14 (0.38)	n=4
4B	K246M(259)	0.69	0.83 (0.05)	5	0.83 (0.06)	0.76 (0.05)	0.95 (0.09)	n=3
5B	K248M(261)	0.79	0.95 (0.06)	4	0.89 (0.09)	0.83 (0.04)	1.01 (0.23)	n=3
8	I253A(266)	< 0.10	0.96 (0.05)	4	1.14 (0.02)	1.18 (0.06)	1.08 (0.14)	n=3
13	T260A(273)	1.09	0.93 (0.09)	4	0.89 (0.14)	0.87 (0.10)	0.89 (0.08)	n=4
20	K274A(287)	1.18	1.02 (0.04)	4	0.86 (0.09)	0.96 (0.10)	1.11 (0.08)	n=3
24	V282A(299)	1.13 (0.07)	2 0.96 (0.02)	4	1.15 (0.13)	1.15 (0.20)	1.00 (0.18)	n=4
28	K288A(305)	0.38 (0.12)	5 0.88 (0.15)	15	1.15 (0.26)	1.14 (0.20)	1.06 (0.04)	n=4
37	Y300F(319)	0.74 (0.10)	2 1.07 (0.15)	4	1.11 (0.04)	1.09 (0.09)	1.01 (0.10)	n=3
43	Q311A(330)	1.62 (0.25)	4 0.93 (0.05)	4	1.11 (0.06)	1.19 (0.13)	0.93 (0.17)	n=3
46	K317A(336)	1.44 (0.18)	4 0.92 (0.17)	6	1.13 (0.05)	1.18 (0.27)	1.10 (0.23)	n=4
47	E318A(337)	0.85	0.92 (0.07)	4	1.04 (0.10)	1.17 (0.23)	1.01 (0.05)	n=3
53	A330Q(349)	0.76	0.96 (0.10)	4	1.01 (0.12)	1.02 (0.02)	0.75 (0.18)	n=3

TABLE 7
CH3 DOMAIN VARIANTS

IG2	Res#EU (Kabat)	FcRn mean sd	n	FcYRI mean sd	n	FcYRIIA mean sd	FcYRIIB mean sd	FcYRIIIA mean sd	n
B1	K338(358)A	1.14		0.90 (0.05)	3	0.78 (0.09)	0.63 (0.08)	0.15 (0.01)	n=2
B1A	K338(358)M	0.78		0.99 (0.08)	3	0.99 (0.13)	0.93 (0.15)	0.49 (0.04)	n=2
B2	K340(360)A	1.02		1.04 (0.07)	3	1.05 (0.18)	0.96 (0.20)	0.84 (0.11)	n=2
B2A	K340(360)M	1.20		1.17 (0.11)	3	1.10 (0.12)	1.20 (0.19)	0.75 (0.12)	n=2
B3	Q342(363)A	1.09		1.13 (0.11)	3	1.01 (0.10)	1.09 (0.23)	0.98 (0.10)	n=2
B4	R344(365)A	0.77		1.04 (0.08)	3	0.89 (0.14)	0.91 (0.04)	0.97 (0.07)	n=4
B5	E345(366)A	1.18		1.06 (0.05)	3	1.03 (0.10)	0.98 (0.10)	0.97 (0.13)	n=4
B6	Q347(368)A	0.95		1.04 (0.06)	3	1.00 (0.03)	0.92 (0.02)	1.04 (0.12)	n=4
B7	R355(376)A	1.06		1.09 (0.07)	3	0.84 (0.09)	0.87 (0.11)	0.98 (0.09)	n=4
B8	E356(377)A	1.21 (0.11)	2	1.05 (0.04)	3	0.90 (0.02)	0.99 (0.13)	0.92 (0.03)	n=3
B9	M358(381)A	0.96		1.06 (0.07)	3	1.11 (0.06)	1.16 (0.25)	0.91 (0.09)	n=3
B10	T359(382)	1.04		1.04 (0.05)	3	1.13 (0.10)	1.15 (0.04)	1.23 (0.26)	n=3
B11	K360(383)A	1.30 (0.08)	4	1.02 (0.04)	3	1.12 (0.10)	1.12 (0.08)	1.23 (0.16)	n=6
B12	N361(384)A	1.16		1.00 (0.03)	3	0.82 (0.07)	0.82 (0.12)	1.08 (0.06)	n=3
B13	Q362(385)A	1.25 (0.24)	3	1.00 (0.04)	3	1.03 (0.10)	1.02 (0.03)	1.03 (0.16)	n=4
B14	Y373(396)A	0.86		0.98 (0.07)	3	0.84 (0.11)	0.75 (0.08)	0.67 (0.04)	n=5
B15	S375(398)A	1.17 (0.19)	5	0.95 (0.02)	3	1.08 (0.06)	1.14 (0.11)	1.04 (0.05)	n=6
B16	D376(399)A	1.45 (0.36)	4	1.00 (0.05)	3	0.80 (0.16)	0.68 (0.14)	0.55 (0.10)	n=5

B17 A378(401)Q	1.32	(0.13)	3	1.06	(0.05)	3	1.40	(0.17)	1.45	(0.17)	1.19	(0.17)	n=5
B18 E380(405)A	2.19	(0.29)	6	1.04	(0.06)	3	1.18	(0.01)	1.07	(0.05)	0.92	(0.12)	n=2
B19 E382(407)A	1.51	(0.18)	4	1.06	(0.03)	3	0.95	(0.11)	0.84	(0.04)	0.76	(0.17)	n=3
B20 S383(408)A	0.74			1.03	(0.03)	3	0.92	(0.04)	0.94	(0.05)	0.88	(0.07)	n=3
B21 N384(410)A	0.88			1.00	(0.01)	3	1.05	(0.19)	1.10	(0.18)	0.96	(0.18)	n=8
B22 Q386(414)A	0.70	(0.10)	2	1.14	(0.08)	3	1.08	(0.13)	1.19	(0.25)	0.98	(0.14)	n=9
B23 E388(416)A	0.64	(0.12)	2	1.15	(0.09)	3	0.87	(0.03)	0.94	(0.09)	0.62	(0.04)	n=3
B24 N389(417)A	0.73			1.00	(0.02)	3	0.98	(0.15)	0.81	(0.04)	0.75	(0.02)	n=3
B25 N390(418)A	0.87			1.06	(0.04)	3	0.99	(0.10)	0.94	(0.02)	0.87	(0.09)	n=3
B26A Y391(419)A	1.14			1.00	(0.08)	3	0.97	(0.10)	0.94	(0.02)	0.86	(0.05)	n=3
B26B Y391(419)F	0.81	(0.10)	2	1.00	(0.01)	3	1.05	(0.12)	1.11	(0.08)	1.01	(0.15)	n=5
B27 K392(420)A	0.97			1.01	(0.08)	3	0.92	(0.20)	0.94	(0.01)	0.79	(0.22)	n=3
B28 L398(426)A	0.94	(0.04)	2	1.13	(0.15)	6	1.17	(0.11)	1.20	(0.08)	0.94	(0.04)	n=3
B29 S400(428)A	0.64	(0.07)	3	1.10	(0.09)	3	0.95	(0.04)	0.99	(0.08)	0.83	(0.07)	n=2
B30 D401(430)A	1.10	(0.09)	3	1.13	(0.16)	6	1.11	(0.12)	1.19	(0.11)	0.97	(0.10)	n=5
B31 D413(444)A	1.21	(0.07)	2	1.00	(0.01)	3	0.83	(0.08)	0.84	(0.06)	0.90	(0.16)	n=2
B32 K414(445)A	1.02			1.00	(0.04)	3	0.64	(0.15)	0.58	(0.18)	0.82	(0.27)	n=3
B33 S415(446)A	0.44			1.04	(0.03)	3	0.90	(0.11)	0.88	(0.05)	0.86	(0.18)	n=2
B34 R416(447)A	1.08			0.96	(0.04)	3	0.68	(0.05)	0.80	(0.05)	0.71	(0.08)	n=2
B35 Q418(449)A	0.77	(0.03)	2	0.98	(0.01)	3	1.00	(0.01)	0.96	(0.02)	0.96	(0.05)	n=2
B36 Q419(450)A	0.76	(0.01)	2	0.97	(0.02)	3	0.68	(0.09)	0.63	(0.07)	0.86	(0.08)	n=3

B37	N421(452)A	0.98		0.99	(0.01)	3	0.90	(0.03)	0.81	(0.0)	0.87	(0.12)	n=2	
B38	V422(453)A	1.01		0.98	(0.02)	3	0.89	(0.0)	0.83	(0.05)	0.83	(0.12)	n=2	
B39	S424(455)A	1.41	(0.14)	3	0.98	(0.03)	3	1.04	(0.06)	1.02	(0.02)	0.88	(0.09)	n=2
B40	E430(461)A	0.93	(0.03)	2	1.05	(0.02)	3	1.24	(0.11)	1.28	(0.10)	1.20	(0.18)	n=5
B41	H433(464)A	0.41	(0.14)	2	0.98	(0.03)	3	0.92	(0.18)	0.79	(0.18)	1.02	(0.15)	n=3
B42	N434(465)A	3.46	(0.37)	7	1.00	(0.04)	3	0.97	(0.07)	0.98	(0.13)	0.74	(0.12)	n=5
B43	H435(466)A	< 0.10		4	1.25	(0.09)	3	0.77	(0.05)	0.72	(0.05)	0.78	(0.03)	n=3
B44	Y436(467)A	< 0.10		2	0.99	(0.02)	2	0.93	(0.05)	0.91	(0.06)	0.91	(0.15)	n=3
B45	T437(468)A	0.99	(0.07)		1.00	(0.02)	3	1.12	(0.18)	1.00	(0.22)	0.77	(0.19)	n=5
B46	Q438(469)A	0.79	(0.05)	2	1.02	(0.05)	3	0.80	(0.10)	0.72	(0.16)	1.01	(0.17)	n=5
B47	K439(470)A	0.70	(0.04)	2	0.98	(0.04)	3	0.78	(0.16)	0.68	(0.22)	0.86	(0.19)	n=4
B48	S440(471)A	0.99			1.01	(0.02)	3	1.10	(0.15)	1.11	(0.26)	0.93	(0.01)	n=3
B49	S442(473)A	0.86			1.02	(0.02)	3	0.98	(0.08)	0.91	(0.11)	0.95	(0.10)	n=5
B50	S444(475)A	0.80			1.01	(0.02)	3	1.07	(0.03)	1.03	(0.03)	0.88	(0.12)	n=2
B51	K447(478)A	0.62	(0.12)	3	1.02	(0.03)	3	0.95	(0.05)	0.91	(0.05)	0.84	(0.09)	n=2

Variants with increased binding to a FcγR generally had binding values ≥ 1.20 as determined in this Example and those with reduced binding to a FcγR generally had binding values ≤ 0.80 as determined in this Example. Variants with increased binding to FcRn generally had binding values ≥ 1.30 as determined in this Example and those with reduced binding to FcRn generally had binding values ≤ 0.70 as determined in this Example.

Aside from alanine variants, various non-alanine substitution variants were made, and the FcR binding activity of those variants is summarized in the following table.

TABLE 8
NON-ALANINE VARIANTS

IG2	Res#EU (Kabat)	FcRn mean sd n	FcγRI mean sd n	FcγRIIA mean sd n	FcγRIIB mean sd n	FcγRIIA mean sd n
222	D249(262)E			0.97	0.99	0.84 n=1
176	T256(269)G			1.10 (0.03)	1.06 (0.07)	0.96 (0.27) n=2
254	T256(269)N			1.03	0.89	1.13 n=1
157	D265(278)N			0.02 (0.01)	0.03 (0.01)	0.02 (0.01) n=3
158	D265(278)E			0.11 (0.04)	0.03 (0.01)	0.02 (0.01) n=3
189	S267(280)G		R131 1.21 (0.05) H131 0.59 (0.09)	0.97 (0.16)	0.09 (0.02)	n=3 n=3
84	H268(281)N			1.33	1.41	0.56 n=1
85	H268(281)S			1.35	1.38	0.81 n=1
87	H268(281)Y			1.19	1.29	0.76 n=1
168	E269(282)D			0.89 (0.10)	0.73 (0.07)	1.13 (0.21) n=2
169	E269(282)Q			0.08 (0.01)	0.16 (0.00)	0.28 (0.03) n=2
92	D270(283)N			0.06 (0.01)	0.10 (0.02)	0.04 (0.00) n=2
93	D270(283)E			0.55 (0.05)	0.38 (0.05)	1.17 (0.01) n=2
223	E272(285)Q			1.93	1.81	0.82 n=1
224	E272(285)N			0.43	0.23	0.50 n=1
167	K274(287)Q			0.86	0.94	0.62 n=1
165	N276(289)K			0.81	0.77	0.61 n=1

233	N276(289)Q	1.09	0.79	0.91	n=1
79	D280(295)N	1.26 (0.07)	1.38 (0.04)	1.13 (0.13)	n=2
149	D280(295)S	1.07 (0.06)	1.04 (0.08)	1.09 (0.06)	n=2
226	E283(300)Q	1.12	1.24	1.19	n=1
227	E283(300)S	1.03	1.07	0.85	n=1
228	E283(300)N	1.18	1.28	0.94	n=1
229	E283(300)D	1.14	1.23	0.95	n=1
23	N286(303)Q	1.52	1.13	0.96	n=1
237	N286(303)S	1.72	1.38	1.32	n=1
238	N286(303)D	1.41	1.23	0.98	n=1
73	K290(307)Q	1.17	1.26	1.40	n=1
75	K290(307)S	1.27	1.34	1.26	n=1
77	K290(307)E	1.10	1.20	1.30	n=1
78	K290(307)R	1.05	1.15	1.08	n=1
177	K290(307)G	1.07	1.21	1.23	n=1
80	R292(309)K	0.71 (0.17)	0.75 (0.10)	1.15 (0.18)	n=3
81	R292(309)H	0.21 (0.09)	0.12 (0.01)	0.92 (0.08)	n=2
82	R292(309)Q	0.47 (0.12)	0.25 (0.06)	0.45 (0.09)	n=3
83	R292(309)N	0.54 (0.16)	0.29 (0.07)	0.88 (0.02)	n=3
144	E293(310)Q	0.85 (0.03)	0.77 (0.13)	0.99 (0.04)	n=2
145	E293(310)D	0.90 (0.02)	0.88 (0.07)	0.37 (0.07)	n=2
147	E293(310)K	1.13 (0.04)	1.31 (0.17)	0.72 (0.08)	n=4

173	E294(311)Q	1.01	0.95	0.84	n=1
174	E294(311)D	0.37	0.26	0.14	n=1
185	Y296(313)H	0.90	0.81	0.92	n=1
186	Y296(313)W	0.96	0.93	1.38	n=1
70	S298(317)G	0.87 (0.17)	0.63 (0.33)	0.46 (0.09)	n=4
71	S298(317)T	0.41 (0.21)	0.40 (0.19)	0.89 (0.20)	n=3
72	S298(317)N	0.08 (0.01)	0.16 (0.03)	0.06 (0.01)	n=2
218	S298(317)V	0.11 (0.06)	0.17 (0.01)	0.33 (0.19)	n=3
219	S298(317)L	1.14 (0.12)	1.42 (0.31)	0.34 (0.04)	n=3
150	V303(322)L	0.89 (0.05)	0.73 (0.10)	0.76 (0.09)	n=4
151	V303(322)T	0.64 (0.11)	0.34 (0.05)	0.20 (0.05)	n=4
217	E318(337)K	1.03	1.08	0.72	n=1
172	K320(339)R	0.71	0.66	0.68	n=1
202	K320(339)M	1.34	1.40	1.27	n=1
204	K320(339)Q	1.23	1.12	1.17	n=1
205	K320(339)E	1.29	1.34	1.12	n=1
235	K320(339)R	1.24	0.95	0.86	n=1
155	K322(341)R	0.87 (0.07)	0.87 (0.21)	0.92 (0.15)	n=3
156	K322(341)Q	0.87 (0.02)	0.92 (0.23)	0.78 (0.18)	n=3
206	K322(341)E	1.38	1.34	0.81	n=1
207	K322(341)N	0.57	0.36	0.04	n=1

213	S324(343)N	1.15	1.09	0.97	n=1
214	S324(343)Q	0.82	0.83	0.78	n=1
215	S324(343)K	0.66	0.37	0.77	n=1
216	S324(343)E	0.82	0.73	0.81	n=1
208	K326(345)S	1.44	1.62	1.37	n=1
209	K326(345)N	1.04	1.00	1.27	n=1
210	K326(345)Q	1.36	1.41	1.15	n=1
211	K326(345)D	1.68	2.01	1.36	n=1
212	K326(345)E	1.34 (0.27)	1.47 (0.33)	1.26 (0.04)	n=2
131	A327(346)S	0.23 (0.06)	0.22 (0.05)	0.06 (0.01)	n=4
159	A327(346)G	0.92 (0.09)	0.83 (0.10)	0.36 (0.05)	n=3
196	A330(349)D	0.18	0.08	0.07	n=1
197	A330(349)K	1.28	1.25	1.28	n=1
198	P331(350)S	1.00	0.86	0.86	n=1
199	P331(350)N	0.86	0.65	0.23	n=1
200	P331(350)E	1.06	0.91	0.42	n=1
203	P331(350)K	0.94	0.71	0.33	n=1
141	E333(352)Q	0.70 (0.05)	0.64 (0.09)	1.10 (0.03)	n=2
142	E333(352)N	0.59 (0.04)	0.52 (0.07)	0.56 (0.10)	n=2
143	E333(352)S			0.94	n=1
152	E333(352)K			0.85 (0.14)	n=3
153	E333(352)R	0.75 (0.04)	0.66 (0.03)	0.84 (0.05)	n=2

154 E333(352)D			1.26 (0.04)	n=3
178 E333(352)G	0.87	0.76	1.05	n=1
179 K334(353)G	0.76 (0.08)	0.60 (0.13)	0.88 (0.22)	n=5
135 K334(353)R	1.15 (0.09)	1.33 (0.18)	0.68 (0.07)	n=5
136 K334(353)Q	1.08 (0.11)	1.10 (0.21)	1.31 (0.26)	n=7
137 K334(353)N	1.16 (0.11)	1.29 (0.30)	1.15 (0.16)	n=7
138 K334(353)S	1.01 (0.11)	1.03 (0.05)	1.19 (0.08)	n=3
139 K334(353)E	0.74 (0.15)	0.72 (0.12)	1.30 (0.09)	n=4
140 K334(353)D	0.51 (0.09)	0.40 (0.03)	1.13 (0.09)	n=4
190 K334(353)M	1.18	1.06	1.01	1.35
191 K334(353)Y	1.15	1.08	1.05	1.31
192 K334(353)W	1.16	0.94	0.91	1.07
193 K334(353)H	1.11	1.09	1.07	1.26
220 K334(353)V		1.13 (0.11)	1.09 (0.15)	1.34 (0.18)
221 K334(353)L		1.05	1.09	1.38
171 T335(354)Q		0.86	0.79	0.84
194 T335(354)E		1.24	1.30	1.19
195 T335(354)K		1.19	1.14	1.30
273 A339(359)T	1.23	1.11	1.23	1.42

The following table summarizes the FcR binding activity of various combination variants.

TABLE 9
COMBINATION VARIANTS

IG2	Res#EU (Kabat)	FcRn mean sd n	FcγRI mean sd n	FcγRIIA mean sd n	FcγRIIB mean sd n	FcγRIIIA mean sd n
96	S267(280)A H268(281)A		1.41	1.72	0.84	n=1
134	E333(352)A K334(353)A		0.72 (0.08)	0.63 (0.13)	1.30 (0.12)	n=5
1059	T256(269)A S298(317)A		0.44 (0.03)	0.22 (0.04)	1.41 (0.06)	n=2
1051	T256(269)A D280(295)A S298(317)A T307(326)A		0.47 (0.01)	0.30 (0.03)	1.21 (0.26)	n=2
106	T256(269)A D280(295)A R292(309)A S298(317)A T307(326)A		0.11	0.08	0.90	n=1
107	S298(317)A E333(352)A		0.34 (0.05)	0.16 (0.08)	1.53 (0.24)	n=5
109	S298(317)A K334(353)A		0.41 (0.07)	0.19 (0.08)	1.62 (0.34)	n=6
110	S298(317)A E333(352)A K334(353)A		0.35 (0.13)	0.18 (0.08)	1.66 (0.42)	n=11
246	S267(280)A E258(271)A		1.62 (0.15)	2.01 (0.45)	1.04 (0.12)	n=2
247	S267(280)A R255(268)A		1.60 (0.18)	1.72 (0.13)	0.88 (0.07)	n=3

248	S267(280)A D280(295)A	1.54 (0.08)	1.96 (0.37)	1.13 (0.07)	n=2
250	S267(280)A E272(285)A	1.51 (0.13)	1.82 (0.32)	0.95 (0.05)	n=3
251	S267(280)A E293(310)A	1.67 (0.11)	1.85 (0.10)	0.92 (0.09)	n=3
264	S267(280)A E258(271)A D280(295)A R255(268)A	1.48 (0.12)	2.03 (0.30)	0.89 (0.04)	n=2
269	E380(405)A N434(465)A	1.02 (0.07)	1.05 (0.11)	1.02	n=2
270	E380(405)A N434(465)A T307(326)A	0.99 (0.06)	0.99 (0.11)	0.96	n=2
271	E380(405)A L309(328)A	0.98	1.04	0.92	n=1
272	N434(465)A K288(305)A	0.94 (0.11)	0.96 (0.17)	0.88	n=2

DISCUSSION

This study includes a complete mapping of human IgG1 for human Fc γ R1, Fc γ R1IA, Fc γ R1IB, Fc γ R1IIA, and FcRn. An alanine-scan of all amino acids in human IgG1 Fc (CH2 and CH3 domains) exposed to solvent, based on the crystal structure of human Fc (Deisenhofer, *Biochemistry* 20:2361-2370 (1981)), was performed. Each exposed amino acid in CH2 and CH3 was individually changed to alanine and the variant IgG assayed against all five human receptors; all variants were evaluated using humanized anti-IgE E27 IgG1 as the parent polypeptide. Fc γ R1 and FcRn are high affinity receptors and monomeric IgG could be evaluated in the assays for these two receptors. Fc γ R1IA, Fc γ R1IB and Fc γ R1IIA are low affinity receptors and required use of an immune complex. Hence, an ELISA-type assay was used for Fc γ R1IA, Fc γ R1IB, and Fc γ R1IIA, in which pre-formed hexamers, consisting of three anti-IgE E27 and three IgE molecules were bound to the Fc γ R and either anti-human IgG Fc-HRP or protein G-HRP used as detection reagent. In order to increase binding, these hexamers could be linked into multimers by addition of human VEGF (using anti-VEGF IgE). The hexamers bound to the low affinity Fc γ R significantly better than the IgG monomers; the multimers bound better than the hexamers (Figs. 15A and 15B). The hexameric complexes were used since these provided sufficient binding and required less IgG. Complexes formed using other antibody:antigen combinations are also possible reagents, as long as the antigen contains at least two identical binding sites per molecule for the antibody. As an example, VEGF contains two binding sites per VEGF dimer for anti-VEGF A.4.6.1 (Kim *et al.*, *Growth Factors* 7:53 (1992) and Kim *et al.* *Nature* 362:841 (1993)). VEGF:anti-VEGF multimers also bound to the low affinity Fc γ R1IA and Fc γ R1IIA (Figs. 16A and 16B).

Once the complete alanine-scan was performed, several classes of alanine variants were found. Some variants exhibited reduced binding to all Fc γ R (G14, Fig. 17), while other variants showed reduced binding only to one Fc γ R (G36, Fig. 17), improved binding only to one Fc γ R (G15, G54, G55, Fig. 17), or simultaneous reduction to one Fc γ R with improvement to another (G16, Fig. 17).

Individual alanine variants were also combined in a single variant Fc region; e.g. combining S298(317)A with K334(353)A improved binding to Fc γ R1IIA more than either S298(317)A or K334(353)A alone (Figs. 18A and B; and compare variants 36, 55, and 109 in Tables 6 and 9) (residue numbers in parentheses are those of the EU index as in Kabat). Similarly, combining S298(317)A with E333(352)A improved binding to Fc γ R1IIA more than either S298(317)A or E333(352)A alone (compare variants 36, 54, and 107 in Tables 6 and 9).

Selected IgG variants were also tested for their binding to Fc γ R transfected into mammalian cells. The α -chain extracellular portion of human Fc γ R1IIA was transfected into CHO cells using a GPI-link, whereas for human Fc γ R1IB the full-length receptor was transfected into

CHO cells. For the variants tested, the pattern of binding to the cells was the same as the pattern of binding in the protein:protein (ELISA) assay (Figs. 18A-B and 19A-B).

One application of these variants is to improve the ADCC effector function of an antibody. This can be achieved by modifying Fc region amino acids at one or more residues which would lead to improved binding to Fc γ RIIIA. Improved Fc γ RIIIA binding would lead to improved binding by NK cells, which carry only Fc γ RIIIA and can mediate ADCC. Selected alanine variants which were either reduced in binding to Fc γ RIIIA (variants 17, 18, 34; Table 6), had no effect on Fc γ RIIIA binding (variant 31; Table 6), or had improved binding to Fc γ RIIIA (variants 30, 36; Table 6) were tested in an *in vitro* ADCC assay using human PBMCs as effector cells. Since the target cells were HER2-overexpressing SKBR3 cells, the IgG Fc variants used in this assay were generated by substituting the V_H/V_L domains of anti-IgE E27 with those from anti-HER2 antibody; HERCEPTIN® (humAb4D5-8 in Table 1 of Carter *et al. PNAS (USA)* 89:4285-4289 (1992)). The pattern of ADCC exhibited by the variants correlated well with the pattern of binding to Fc γ RIIIA (Figs. 20 and 21). Notably the variant which showed the best improvement in binding to Fc γ RIIIA in protein:protein assays, variant 36 S298(317)A, also showed improvement in ADCC compared to wildtype HERCEPTIN® at 1.25 ng/ml (Fig. 21).

EXAMPLE 5

Bind of Fc Variants to Polymorphic Fc Receptors

Allelic variants of several of the human Fc γ R have been found in the human population. These allelic variant forms have been shown to exhibit differences in binding of human and murine IgG and a number of association studies have correlated clinical outcomes with the presence of specific allelic forms (reviewed in Lehmbecher *et al. Blood* 94(12):4220-4232 (1999)). Several studies have investigated two forms of Fc γ RIIA, R131 and H131, and their association with clinical outcomes (Hatta *et al. Genes and Immunity* 1:53-60 (1999); Yap *et al. Lupus* 8:305-310 (1999); and Lorenz *et al. European J. Immunogenetics* 22:397-401 (1995)). Two allelic forms of Fc γ RIIIA, F158 and V158, are only now being investigated (Lehmbecher *et al., supra*; and Wu *et al. J. Clin. Invest.* 100(5):1059-1070 (1997)). In this example, selected IgG variants were tested against both allelic forms of Fc γ RIIA or Fc γ RIIIA. Fc receptor binding assays were performed essentially as described in the above examples. However, for Fc γ RIIIA-V158, both (a) the low affinity receptor binding assay of Example 1 (which analyzes binding of the IgG complex to Fc γ RIIIA-V158); and (b) the high affinity Fc γ R binding assay of Example 4 (which analyzes binding of IgG monomer to Fc γ RIIIA-V158) were carried out. The results of these studies are summarized in Table 10 below.

TABLE 10
Binding of Variants to FcγRIIA and FcγRIIA Polymorphic Receptors

IG2	Res#EU (Kabat)	IgG Complex			IgG Complex			IgG Complex			IgG Complex			IgG Monomer			
		FcγRIIA-R131	mean	sd	n	FcγRIIA-H131	mean	sd	n	FcγRIIA-F158	mean	sd	n	FcγRIIA-V158	mean	sd	n
11	T256(269)A	1.41	(0.27)	9					1.32 (0.18)	9	0.97 (0.03)	2	1.20	1			
254	T256(269)N	1.03		1					1.13	1	0.95	1	0.88	1			
14	D265(278)A	0.07	(0.01)	4					0.09 (0.06)	4	0.01	1					
15	S267(280)A	1.64	(0.18)	7					1.05 (0.03)	2	1.14 (0.25)	7					
189	S267(280)G	1.21	(0.05)	3					0.59 (0.09)	3	0.09 (0.02)	3					
16	H268(281)A	1.22	(0.14)	12					1.09 (0.01)	2	0.52 (0.09)	12					
25	E283(300)A	1.24	(0.23)	5					1.01 (0.14)	5			0.78	1			
226	E283(300)Q	1.12		1					1.19	1			0.89	1			
227	E283(300)S	1.03		1					0.85	1			0.83	1			
228	E283(300)N	1.18		1					0.94	1			0.63	1			
229	E283(300)D	1.14		1					0.95	1			0.67	1			
30	K290(307)A	1.29	(0.21)	7					1.28 (0.21)	7	1.12 (0.05)	2	1.13	1			
73	K290(307)Q	1.17		1					1.40	1	1.02	1	1.30	1			
75	K290(307)S	1.27		1					1.26	1	1.05	1	1.62	1			
77	K290(307)E	1.10		1					1.30	1	0.98	1	1.50	1			
78	K290(307)R	1.05		1					1.08	1	1.07	1	1.24	1			
177	K290(307)G	1.07		1					1.23	1	1.11	1	2.29	1			

31	R292(309)A	0.27 (0.14)	9	0.90 (0.18)	9	0.94	1
80	R292(309)K	0.71 (0.17)	3	1.15 (0.18)	3	1.64	1
81	R292(309)H	0.21 (0.09)	2	0.92 (0.08)	2	1.21	1
82	R292(309)Q	0.47 (0.12)	3	0.45 (0.09)	3	0.56	1
83	R292(309)N	0.54 (0.16)	3	0.88 (0.02)	3	0.91	1
144	E293(310)Q	0.85 (0.03)	2	0.99 (0.04)	2	1.00	1
33	E294(311)A	0.87 (0.19)	5	0.66 (0.14)	5	0.68	1
173	E294(311)Q	1.01	1	0.84	1	0.79	1
174	E294(311)D	0.37	1	0.14	1	0.26	1
36	S298(317)A	0.40 (0.08)	12	1.30 (0.18)	12	1.02 (0.04)	2
70	S298(317)G	0.87 (0.17)	4	0.46 (0.09)	4	0.88	1
71	S298(317)T	0.41 (0.21)	3	0.89 (0.20)	3	0.96	1
72	S298(317)N	0.08 (0.01)	2	0.06 (0.01)	2	0.66	1
218	S298(317)V	0.11 (0.06)	3	0.33 (0.19)	3	0.88	1
219	S298(317)L	1.14 (0.12)	3	0.34 (0.04)	3	0.83	1
40	V305(324)A	1.12 (0.12)	4	1.04	1	0.84 (0.15)	4
41	T307(326)A	1.19 (0.37)	12	1.37 (0.13)	2	1.12 (0.18)	12
45	N315(334)A	1.15 (0.06)	5	1.11 (0.06)	2	1.07 (0.21)	5
46	K317(336)A	1.13 (0.05)	4	1.04	1	1.10 (0.23)	4
48	K320(339)A	1.12 (0.11)	4	1.16	1	0.87 (0.17)	4
54	E333(352)A	0.92 (0.12)	10	1.27 (0.17)	10	1.10 (0.10)	2
						1.29	1

141	E333(352)Q	0.70 (0.05) 2	1.10 (0.03) 2	1.05	1	1.00	1
142	E333(352)N	0.59 (0.04) 2	0.56 (0.10) 2	0.64	1	0.56	1
143	E333(352)S		0.94	1	0.99	1	1.07
152	E333(352)K		0.85 (0.14) 3	0.88	1	0.81	1
153	E333(352)R	0.75 (0.04) 2	0.84 (0.05) 2	0.93	1	0.83	1
154	E333(352)D		1.26 (0.04) 3	1.00	1	1.70	1
178	E333(352)G	0.87	1.05	1		1.23	1
55	K334(353)A	1.01 (0.15) 17	1.39 (0.19) 17	1.07 (0.09) 3	1.60 (0.01) 2		
135	K334(353)R	1.15 (0.09) 5	0.68 (0.07) 5		0.88	1	
136	K334(353)Q	1.08 (0.11) 7	1.31 (0.26) 7	1.27 (0.01) 2	1.92	1	
137	K334(353)N	1.16 (0.11) 7	1.15 (0.16) 7	1.19 (0.06) 2	1.70	1	
138	K334(353)S	1.01 (0.11) 3	1.19 (0.08) 3	1.25	1	1.82	1
139	K334(353)E	0.74 (0.15) 4	1.30 (0.09) 4	1.17	1	2.75	1
140	K334(353)D	0.51 (0.09) 4	1.13 (0.09) 4	1.07	1		
179	K334(353)G	0.76 (0.08) 5	0.88 (0.22) 5	0.94	1	1.28	1
190	K334(353)M	1.06	1.35	1	0.99	1	2.08
191	K334(353)Y	1.08	1.31	1	0.98	1	1.72
192	K334(353)W	0.94	1.07	1	0.96	1	1.53
193	K334(353)H	1.09	1.26	1	0.97	1	2.06
220	K334(353)V	1.13 (0.11) 3	1.34 (0.18) 3	1.00	1	2.89	1
221	K334(353)L	1.05	1.38	1	0.96	1	3.59

65	P331(350)A	1.29 (0.14)	3	1.03 (0.19)	3	0.96	1	0.78	1
198	P331(350)S	1.00	1	0.86	1			0.54	1
199	P331(350)N	0.86	1	0.23	1			0.24	1
200	P331(350)E	1.06	1	0.42	1			0.36	1
203	P331(350)K	0.94	1	0.33	1			0.26	1
96	S267(280)A H268(281)A	1.54 (0.12)	3	1.07 (0.06)	2	0.84	1		
110	S298(317)A E333(352)A K334(353)A	0.35 (0.13)	11	1.66 (0.42)	11	1.19 (0.18)	3		
271	E380(405)A L309(328)A	0.98	1	0.92	1			1.10	1

For FcγRIIIA, the pattern of binding of the selected IgG1 variants to the relatively higher affinity FcγRIIIA-V158 was the same as for the relatively lower affinity FcγRIIIA-F158 (the F158 form was used in assaying all variants). IgG1 variants which showed improved binding to the FcγRIIIA-F158 form also showed improved binding to the FcγRIIIA-V158 form though the improvement was not as pronounced. For FcγRIIA-R131 (used in assaying all variants) and FcγRIIA-H131, the binding pattern of the selected IgG1 variants did show some distinct differences. S267(280)A, H268(281)A, and S267(280)A/H268(281)A exhibited improved binding to FcγRIIA-R131, compared to native IgG1, but not to FcγRIIA-H131. In contrast, S267(280)G showed improved binding to FcγRIIA-R131 but reduced binding to FcγRIIA-H131 (Table 10). Other variants bound similarly to both allelic FcγRIIA forms: V305(324)A, T307(326)A, N315(324)A, K317(336)A, and K320(339)A.

WHAT IS CLAIMED IS:

1. A variant of a parent polypeptide comprising an Fc region, which variant mediates antibody-dependent cell-mediated cytotoxicity (ADCC) in the presence of human effector cells more effectively, or binds an Fc gamma receptor (FcγR) with better affinity, than the parent polypeptide and comprises at least one amino acid modification in the Fc region.
2. The variant of claim 1 which comprises an antibody.
3. The variant of claim 1 wherein the parent polypeptide Fc region comprises a human IgG Fc region.
4. The variant of claim 3 wherein the human IgG Fc region comprises a human IgG1, IgG2, IgG3 or IgG4 Fc region.
5. The variant of claim 1 which mediates ADCC about 1.5 fold to about 100 fold more effectively than the parent polypeptide.
6. The variant of claim 1 which binds an FcγRIII with better affinity than the parent polypeptide.
7. The variant of claim 6 which further binds an FcγRII with worse affinity than the parent polypeptide.
8. The variant of claim 1 which comprises at least one amino acid substitution in the Fc region.
9. The variant of claim 1 which comprises at least one amino acid modification in a CH2 domain of the Fc region.
10. The variant of claim 1 which comprises at least one amino acid modification in the Fc region, other than in a lower hinge region thereof.
11. The variant of claim 1 which comprises an amino acid substitution at any one or more of amino acid positions 256, 290, 298, 312, 326, 330, 333, 334, 360, 378 or 430 of the Fc region, wherein the numbering of the residues in the Fc region is that of the EU index as in Kabat.
12. The variant of claim 11 which comprises two or more amino acid substitutions at the amino acid positions listed therein.
13. The variant of claim 11 which comprises three or more amino acid substitutions at the amino acid positions listed therein.
14. A polypeptide comprising a variant Fc region with altered Fc gamma receptor (FcγR) binding affinity, which polypeptide comprises an amino acid modification at any one or more of amino acid positions 238, 239, 248, 249, 252, 254, 255, 256, 258, 265, 267, 268, 269, 270, 272, 276, 278, 280, 283, 285, 286, 289, 290, 292, 293, 294, 295, 296, 298, 301, 303, 305, 307, 309, 312, 315, 320, 322, 324, 326, 327, 329, 330, 331, 333, 334, 335, 337, 338, 340, 360, 373, 376, 378, 382, 388, 389, 398, 414, 416, 419, 430, 434, 435, 437, 438 or 439 of the Fc region, wherein the numbering of the residues in the Fc region is that of the EU index as in Kabat.

15. The polypeptide of claim 14 wherein the variant Fc region comprises a variant human IgG Fc region.

16. The polypeptide of claim 14 which displays reduced binding to an FcγR and comprises an amino acid modification at any one or more of amino acid positions 238, 239, 248,
5 249, 252, 254, 265, 268, 269, 270, 272, 278, 289, 292, 293, 294, 295, 296, 298, 301, 303, 322, 324, 327, 329, 333, 335, 338, 340, 373, 376, 382, 388, 389, 414, 416, 419, 434, 435, 437, 438 or 439 of the Fc region, wherein the numbering of the residues in the Fc region is that of the EU index as in Kabat.

17. The polypeptide of claim 14 which displays reduced binding to an FcγRI.

10 18. The polypeptide of claim 17 which displays reduced binding to the FcγRI and comprises an amino acid modification at any one or more of amino acid 238, 265, 269, 270, 327 or 329 of the Fc region, wherein the numbering of the residues in the Fc region is that of the EU index as in Kabat.

19. The polypeptide of claim 14 which displays reduced binding to an FcγRII.

15 20. The polypeptide of claim 19 which displays reduced binding to the FcγRII and comprises an amino acid modification at any one or more of amino acid positions 238, 265, 269, 270, 292, 294, 295, 298, 303, 324, 327, 329, 333, 335, 338, 373, 376, 414, 416, 419, 435, 438 or 439 of the Fc region, wherein the numbering of the residues in the Fc region is that of the EU index as in Kabat.

20 21. The polypeptide of claim 14 which displays reduced binding to an FcγRIII.

22. The polypeptide of claim 21 which displays reduced binding to the FcγRIII and comprises an amino acid modification at any one or more of amino acid 238, 239, 248, 249, 252, 254, 265, 268, 269, 270, 272, 278, 289, 293, 294, 295, 296, 301, 303, 322, 327, 329, 338, 340, 373, 376, 382, 388, 389, 416, 434, 435 or 437 of the Fc region, wherein the numbering of the
25 residues in the Fc region is that of the EU index as in Kabat.

23. The polypeptide of claim 14 which displays increased binding to an FcγR and comprises an amino acid modification at any one or more of amino acid positions 255, 256, 258, 267, 268, 272, 276, 280, 283, 285, 286, 290, 298, 301, 305, 307, 309, 312, 315, 320, 322, 326, 330, 331, 333, 334, 337, 340, 360, 378, 398 or 430 of the Fc region, wherein the numbering of the
30 residues in the Fc region is that of the EU index as in Kabat.

24. The polypeptide of claim 23 which displays increased binding to an FcγRIII.

25. The polypeptide of claim 24 which further displays decreased binding to an FcγRII.

26. The polypeptide of claim 25 which displays increased binding to the FcγRIII and further displays decreased binding to the FcγRII, wherein the polypeptide comprises an amino
35 acid modification at positions 298 and/or 333 of the Fc region, wherein the numbering of the residues in the Fc region is that of the EU index as in Kabat.

27. The polypeptide of claim 23 which displays increased binding to an FcγRII.
28. The polypeptide of claim 27 which displays increased binding to the FcγRII and comprises an amino acid modification at any one or more of amino acid 255, 256, 258, 267, 268, 272, 276, 280, 283, 285, 286, 290, 301, 305, 307, 309, 312, 315, 320, 322, 326, 330, 331, 337, 340, 378, 398 or 430 of the Fc region, wherein the numbering of the residues in the Fc region is that of the EU index as in Kabat.
29. The polypeptide of claim 27 which further displays decreased binding to an FcγRIII.
30. The polypeptide of claim 29 which displays increased binding to the FcγRII and further displays decreased binding to the FcγRIII, wherein the polypeptide comprises an amino acid modification at any one or more of amino acid positions 268, 272, 298, 301, 322 or 340 of the Fc region, wherein the numbering of the residues in the Fc region is that of the EU index as in Kabat.
31. A polypeptide comprising a variant Fc region with altered neonatal Fc receptor (FcRn) binding affinity, which polypeptide comprises an amino acid modification at any one or more of amino acid positions 238, 252, 253, 254, 255, 256, 265, 272, 286, 288, 303, 305, 307, 309, 311, 312, 317, 340, 356, 360, 362, 376, 378, 380, 382, 386, 388, 400, 413, 415, 424, 433, 434, 435, 436, 439 or 447 of the Fc region, wherein the numbering of the residues in the Fc region is that of the EU index as in Kabat.
32. The polypeptide of claim 31 which displays reduced binding to an FcRn.
33. The polypeptide of claim 32 which displays reduced binding to the FcRn and comprises an amino acid modification at any one or more of amino acid positions 252, 253, 254, 255, 288, 309, 386, 388, 400, 415, 433, 435, 436, 439 or 447 of the Fc region, wherein the numbering of the residues in the Fc region is that of the EU index as in Kabat.
34. The polypeptide of claim 31 which displays increased binding to FcRn.
35. The polypeptide of claim 34 which displays increased binding to FcRn and comprises an amino acid modification at any one or more of amino acid positions 238, 256, 265, 272, 286, 303, 305, 307, 311, 312, 317, 340, 356, 360, 362, 376, 378, 380, 382, 413, 424 or 434 of the Fc region, wherein the numbering of the residues in the Fc region is that of the EU index as in Kabat.
36. A composition comprising the polypeptide variant of claim 1 and a pharmaceutically acceptable carrier.
37. The composition of claim 36 which is sterile.
38. Isolated nucleic acid encoding the polypeptide variant of claim 1.
39. A vector comprising the nucleic acid of claim 38.
40. A host cell containing the vector of claim 39.

41. A method for producing a polypeptide variant comprising culturing the host cell of claim 40 so that the nucleic acid is expressed.

42. The process of claim 41 further comprising recovering the polypeptide variant from the host cell culture.

5 43. A method for treating a disorder in a mammal comprising administering to the mammal a therapeutically effective amount of the polypeptide variant of claim 1.

44. A method for making a variant Fc region with altered Fc receptor (FcR) binding affinity, or altered antibody-dependent cell-mediated cytotoxicity (ADCC) activity, comprising:

10 (a) introducing one or more amino acid modifications into an Fc region of a parent polypeptide in order to generate a variant Fc region;

(b) determining binding of the variant Fc region to an FcR, or determining ADCC activity of the variant Fc region.

45. The method of claim 44 wherein step (b) comprises determining binding of the variant Fc region to an FcR *in vitro*.

15 46. The method of claim 44 wherein step (b) comprises identifying a variant Fc region with improved FcR binding affinity, or with improved ADCC activity.

47. The method of claim 44 wherein the FcR is human Fc gamma receptor III (FcγRIII).

48. The method of claim 44 wherein step (b) comprises determining binding of the variant Fc region to at least two different FcRs.

20 49. The method of claim 48 wherein the Fc receptors include human Fc gamma receptor II (FcγRII) and human Fc gamma receptor III (FcγRIII).




FIG. 1




FIG. 2




FIG. 3

FIG. 4A (E27) - Light Chain

DIQLTQSPSS LSASVGDRVT ITCRASKPVD GEGDSYMNWY QOKPGKAPKL LIYAASYLES GVPSRFSGSG
SGTDFTLTIS SLPEDFATY YCQQSHEDPY TFGQGTKVEI KRTVAAPSVF IFPPSDEQLK SGTASVVCLL
NNFYBREAKV QWKVDNALQS GNSQESVTEQ DSKDSTYSLS STLTLSKADY EKHKVYACEV THQGLSSPVT
KSFNRGEC

FIG. 4B (E27) - Heavy Chain

EVQLVESGGG LVQPGGSLRL SCAVSGYSIT SGYSWNWIRQ APKGLEWVA SIKYGETKY NPSVKGRITI
SRDDSKNTFY LQMNSLRAED TAVYVCARGS HYFGHWHFAV WQGTLVTVS SASTKGPSVF PLAPSSKSTS
GGTAALGCLV KDYFPEPTV SWNSGALTSG VHTFPAVLQS SGLYSLSSV TVPSSSLGTQ TYICNVNHP
SNTKVDKKVE PKSCDKTHC PPCPAPELLG GPSVFLFPPK PKDTLMISRT PEVTCVVVDV SHEDPEVKFN
WYVDGVEVHN AKTKPREEQY NSTYRVVSVL TVLHQDWLNG KEYCKVSNK ALPAPIEKTI SKAKGQPREP
QVYTLPPSRE EMTKNQVSLT CLVKGFPYPSD IAVEWESNGQ PENNYKTTTP VLDSDGSFFL YSKLTVDKSR
WQQGNVFSCS VMHEALHNHY TQKSLSLSPG K




Fig. 5




FIG. 6




FIG. 7




FIG. 8




FIG. 9

10/25




FIG. 10

BEST AVAILABLE COPY




FIG. 11

12/25




FIG. 12




FIG. 13

EC50 of wt-C2B8 = 1.54
EC50 of A327g (C2B8) = 1.08






FIG. 14

FcγRIIB



FcγRIIA



FcγRIIA




FIG. 16A

FcγRIIA




FIG. 16B




FIG. 17




FIG. 18A




FIG. 18B




FIG. 19A




FIG. 19B




FIG. 20




FIG. 21

	230	240	250	260	270
humIgG1	PAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYV				
humIgG2	PAP-PVAGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVQFNWYV				
humIgG3	PAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVQFKWYV				
humIgG4	PAPEFLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSEQEDPEVQFNWYV				
murIgG1	---TVPEVSSVFIFPPKPKDVLITLTPKVTCTVVVDISKDDPEVQFSWFV				
murIgG2A	PAPNLLGGPSVFIFFPKIKDVLMLISLSPIVTCVVVDVSEDDPDVQISWFV				
murIgG2B	PAPNLEGGPSVFIFFPNIKDVLMLISLTPKVTCTVVVDVSEDDPDVQISWFV				
murIgG3	PPGNILGGPSVFIFFPKPKDALMISLTPKVTCTVVVDVSEDDPDVHVSWFV				
	280	290	300	310	320
humIgG1	DGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALP				
humIgG2	DGVEVHNAKTKPREEQFNSTFRVSVLTVVHQDWLNGKEYKCKVSNKGLP				
humIgG3	DGVEVHNAKTKPREEQFNSTFRVSVLTVLHQDWLNGKEYKCKVSNKALP				
humIgG4	DGVEVHNAKTKPREEQFNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKGLP				
murIgG1	DDVEVHTAQTQPREEQFNSTFRSVSELPIMHQDCLNGKEFKCRVNSAAFP				
murIgG2A	NNVEVHTAQTQTHREDYNSTLRVVSALPIQHQQDWMSGKEFKCKVNNKDL				
murIgG2B	NNVEVHTAQTQTHREDYNSTIRVVSHPHQDQDWMSGKEFKCKVNNKDL				
murIgG3	DNKEVHTAWTQPREAQYNSTFRVVSALPIQHQQDWMRGKEFKCKVNNKALP				
	330	340	350	360	370
humIgG1	APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAV				
				D L	
humIgG2	APIEKTISKTKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAV				
humIgG3	APIEKTISKTKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAV				
humIgG4	SSIEKTISKAKGQPREPQVYTLPPSQEEMTKNQVSLTCLVKGFYPSDIAV				
murIgG1	APIEKTISKTKGRPKAPQVYTIPPPKEQMAKDKVSLTCMITDFFPEDITV				
murIgG2A	APIERTISKPKGSVRAPQVYVLPPEEEMTKKQVTLTCMVTDFMPEDIYV				
murIgG2B	SPIERTISKPKGLVRAPQVYTLPPPAEQLSRKDVSLTCLVVGFNPGDISV				
murIgG3	APIERTISKPKGRAQTQVYTIPPPREQMSKKKVSILTCLVTNFFSEAISV				
	380	390	400	410	420
humIgG1	EWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFCFSVMH				
humIgG2	EWESNGQPENNYKTTPPMLDSDGSFFLYSKLTVDKSRWQQGNVFCFSVMH				
humIgG3	EWESSGQPENNYNTTPPMLDSDGSFFLYSKLTVDKSRWQQGNIFCFSVMH				
humIgG4	EWZSNGQPENNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFCFSVMH				
murIgG1	EWQWNGQPAENYKNTQPIMDTDGSYFVYSKLVNQKSNWEAGNTFTCSVLH				
murIgG2A	EWTNNGKTELNYKNTPEVLDSDGSYFMYSKLRVEKKNWVERNSYSCSVVH				
murIgG2B	EWTNNGHTEENYKDTAPVLDSDGSYFIYSKLVNMKTSKWEKTDSFSCNVRH				
murIgG3	EWERNGELEQDYKNTPPILDSDGTYFLYSKLTVDTDQSWLQGEIFTCSVVH				
	430	440			
humIgG1	EALHNHYTQKSLSLSPGK				
humIgG2	EALHNHYTQKSLSLSPGK				
humIgG3	EALHNRFTQKSLSLSPGK				
humIgG4	EALHNHYTQKSLSLSLGLK				
murIgG1	EGLHNHHTTEKSLSHSPGK				
murIgG2A	EGLHNHHTTKSFSRTPGK				
murIgG2B	EGLKNYYLKKTISRSPGK				
murIgG3	EALHNHHTQKNLSRSPGK				

FIG. 22A

Percent Identity Among Fc Sequences

	1	2	3	4	5	6	7	8
1. humIgG1	-	94	94	94	64	66	63	68
2. humIgG2		-	93	92	65	63	60	67
3. humIgG3			-	91	64	64	61	67
4. humIgG4				-	62	64	61	64
5. murIgG1					-	65	61	67
6. murIgG2A						-	77	70
7. murIgG2B							-	68
8. murIgG3								-

FIG. 22B

	230	240	250	260	270
humIgG1	PAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYV				
humIgG2	PAP-PVAGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVQFNWYV				
humIgG3	PAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVQFKWYV				
humIgG4	PAPEFLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSQEDPEVQFNWYV				
	****			*	* *
	280	290	300	310	320
humIgG1	DGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALP				
humIgG2	DGVEVHNAKTKPREEQFNSTFRVVSFLTIVVHQDWLNGKEYKCKVSNKGLP				
humIgG3	DGVEVHNAKTKPREEQFNSTFRVVSFLTIVLHQDWLNGKEYKCKVSNKALP				
humIgG4	DGVEVHNAKTKPREEQFNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKGLP				
		*	*	*	*
	330	340	350	360	370
humIgG1	APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAV				
				D L	
humIgG2	APIEKTISKTKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAV				
humIgG3	APIEKTISKTKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAV				
humIgG4	SSIEKTISKAKGQPREPQVYTLPPSQEEMTKNQVSLTCLVKGFYPSDIAV				
	**	*		*	
	380	390	400	410	420
humIgG1	EWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFCFSVMH				
humIgG2	EWESNGQPENNYKTTPPMLDSDGSFFLYSKLTVDKSRWQQGNVFCFSVMH				
humIgG3	EWESSGQPENNYNTTPPMLDSDGSFFLYSKLTVDKSRWQQGNIFCFSVMH				
humIgG4	EWESNGQPENNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFCFSVMH				
	*	*	*	*	* *
	430	440			
humIgG1	EALHNHYTQKSLSLSPGK				
humIgG2	EALHNHYTQKSLSLSPGK				
humIgG3	EALHNRFTQKSLSLSPGK				
humIgG4	EALHNHYTQKSLSLGLK				
	**		*		

FIG. 23




FIG. 24

Sequence Listing

<110> Genentech, Inc.
 <120> Polypeptide Variants with Altered Effector Function
 <130> P1726R1PCT
 5 <141> 2000-01-14
 <150> US 60/116,023
 <151> 1999-01-15
 <160> 11
 <210> 1
 10 <211> 218
 <212> PRT
 <213> Artificial Sequence
 <220>
 <221> Artificial Sequence
 15 <222> 1-218
 <223> Sequence is completely synthesized
 <400> 1
 Asp Ile Gln Leu Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val
 1 5 10 15
 20 Gly Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Lys Pro Val Asp
 20 25 30
 Gly Glu Gly Asp Ser Tyr Met Asn Trp Tyr Gln Gln Lys Pro Gly
 35 40 45
 25 Lys Ala Pro Lys Leu Leu Ile Tyr Ala Ala Ser Tyr Leu Glu Ser
 50 55 60
 Gly Val Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe
 65 70 75
 Thr Leu Thr Ile Ser Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr
 80 85 90
 30 Tyr Cys Gln Gln Ser His Glu Asp Pro Tyr Thr Phe Gly Gln Gly
 95 100 105
 Thr Lys Val Glu Ile Lys Arg Thr Val Ala Ala Pro Ser Val Phe
 110 115 120
 35 Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr Ala Ser
 125 130 135
 Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys Val
 140 145 150
 Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu
 155 160 165
 40 Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser
 170 175 180
 Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val
 185 190 195

Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr
 200 205 210
 Lys Ser Phe Asn Arg Gly Glu Cys
 215 218
 5 <210> 2
 <211> 451
 <212> PRT
 <213> Artificial Sequence
 <220>
 10 <221> Artificial Sequence
 <222> 1-451
 <223> Sequence is completely synthesized
 <400> 2
 15 Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly
 1 5 10 15
 Gly Ser Leu Arg Leu Ser Cys Ala Val Ser Gly Tyr Ser Ile Thr
 20 20 25 30
 Ser Gly Tyr Ser Trp Asn Trp Ile Arg Gln Ala Pro Gly Lys Gly
 35 40 45
 20 Leu Glu Trp Val Ala Ser Ile Lys Tyr Ser Gly Glu Thr Lys Tyr
 50 55 60
 Asn Pro Ser Val Lys Gly Arg Ile Thr Ile Ser Arg Asp Asp Ser
 65 70 75
 25 Lys Asn Thr Phe Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp
 80 85 90
 Thr Ala Val Tyr Tyr Cys Ala Arg Gly Ser His Tyr Phe Gly His
 95 100 105
 Trp His Phe Ala Val Trp Gly Gln Gly Thr Leu Val Thr Val Ser
 110 115 120
 30 Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser
 125 130 135
 Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val
 140 145 150
 35 Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly
 155 160 165
 Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln Ser
 170 175 180
 Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser
 185 190 195
 40 Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro
 200 205 210
 Ser Asn Thr Lys Val Asp Lys Lys Val Glu Pro Lys Ser Cys Asp
 215 220 225
 45 Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly
 230 235 240

Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu
 245 250 255
 Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val
 260 265 270
 5 Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly
 275 280 285
 Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr
 290 295 300
 10 Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln
 305 310 315
 Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys
 320 325 330
 Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly
 335 340 345
 15 Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu
 350 355 360
 Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly
 365 370 375
 20 Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln
 380 385 390
 Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp
 395 400 405
 Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg
 410 415 420
 25 Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala
 425 430 435
 Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly
 440 445 450
 30 Lys
 451
 <210> 3
 <211> 218
 <212> PRT
 <213> homo sapiens
 35 <400> 3
 Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro
 1 5 10 15
 Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val
 20 25 30
 40 Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys
 35 40 45
 Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr
 50 55 60
 Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser

				65					70					75	
	Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu	Tyr
				80						85				90	
5	Lys	Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu	Pro	Ala	Pro	Ile	Glu	Lys
				95						100				105	
	Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr
				110						115				120	
	Thr	Leu	Pro	Pro	Ser	Arg	Glu	Glu	Met	Thr	Lys	Asn	Gln	Val	Ser
				125						130				135	
10	Leu	Thr	Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val
				140						145				150	
	Glu	Trp	Glu	Ser	Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr	Thr
				155						160				165	
15	Pro	Pro	Val	Leu	Asp	Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	Lys
				170						175				180	
	Leu	Thr	Val	Asp	Lys	Ser	Arg	Trp	Gln	Gln	Gly	Asn	Val	Phe	Ser
				185						190				195	
	Cys	Ser	Val	Met	His	Glu	Ala	Leu	His	Asn	His	Tyr	Thr	Gln	Lys
				200						205				210	
20	Ser	Leu	Ser	Leu	Ser	Pro	Gly	Lys							
				215				218							
	<210> 4														
	<211> 218														
	<212> PRT														
25	<213> homo sapiens														
	<400> 4														
	Pro	Ala	Pro	Glu	Leu	Leu	Gly	Gly	Pro	Ser	Val	Phe	Leu	Phe	Pro
	1				5					10				15	
30	Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu	Val
				20						25				30	
	Thr	Cys	Val	Val	Val	Asp	Val	Ser	His	Glu	Asp	Pro	Glu	Val	Lys
				35						40				45	
	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn	Ala	Lys	Thr
				50						55				60	
35	Lys	Pro	Arg	Glu	Glu	Gln	Tyr	Asn	Ser	Thr	Tyr	Arg	Val	Val	Ser
				65						70				75	
	Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu	Tyr
				80						85				90	
40	Lys	Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu	Pro	Ala	Pro	Ile	Glu	Lys
				95						100				105	
	Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr
				110						115				120	
	Thr	Leu	Pro	Pro	Ser	Arg	Asp	Glu	Leu	Thr	Lys	Asn	Gln	Val	Ser
				125						130				135	

Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val
 140 145 150
 Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr
 155 160 165
 5 Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys
 170 175 180
 Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser
 185 190 195
 10 Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys
 200 205 210
 Ser Leu Ser Leu Ser Pro Gly Lys
 215 218
 <210> 5
 <211> 217
 15 <212> PRT
 <213> homo sapiens
 <400> 5
 Pro Ala Pro Pro Val Ala Gly Pro Ser Val Phe Leu Phe Pro Pro
 1 5 10 15
 20 Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr
 20 25 30
 Cys Val Val Val Asp Val Ser His Glu Asp Pro Glu Val Gln Phe
 35 40 45
 25 Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys
 50 55 60
 Pro Arg Glu Glu Gln Phe Asn Ser Thr Phe Arg Val Val Ser Val
 65 70 75
 Leu Thr Val Val His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys
 80 85 90
 30 Cys Lys Val Ser Asn Lys Gly Leu Pro Ala Pro Ile Glu Lys Thr
 95 100 105
 Ile Ser Lys Thr Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr
 110 115 120
 35 Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu
 125 130 135
 Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu
 140 145 150
 Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro
 155 160 165
 40 Pro Met Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu
 170 175 180
 Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys
 185 190 195
 Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser

	200	205	210
	Leu Ser Leu Ser Pro Gly Lys		
	215	217	
	<210> 6		
5	<211> 218		
	<212> PRT		
	<213> homo sapiens		
	<400> 6		
10	Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro		
	1	5	10 15
	Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val		
		20	25 30
	Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro Glu Val Gln		
		35	40 45
15	Phe Lys Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr		
		50	55 60
	Lys Pro Arg Glu Glu Gln Phe Asn Ser Thr Phe Arg Val Val Ser		
		65	70 75
20	Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr		
		80	85 90
	Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys		
		95	100 105
	Thr Ile Ser Lys Thr Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr		
		110	115 120
25	Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser		
		125	130 135
	Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val		
		140	145 150
30	Glu Trp Glu Ser Ser Gly Gln Pro Glu Asn Asn Tyr Asn Thr Thr		
		155	160 165
	Pro Pro Met Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys		
		170	175 180
	Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Ile Phe Ser		
		185	190 195
35	Cys Ser Val Met His Glu Ala Leu His Asn Arg Phe Thr Gln Lys		
		200	205 210
	Ser Leu Ser Leu Ser Pro Gly Lys		
		215	218
	<210> 7		
40	<211> 218		
	<212> PRT		
	<213> homo sapiens		
	<400> 7		
45	Pro Ala Pro Glu Phe Leu Gly Gly Pro Ser Val Phe Leu Phe Pro		
	1	5	10 15

Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val
 20 25 30
 Thr Cys Val Val Val Asp Val Ser Gln Glu Asp Pro Glu Val Gln
 35 40 45
 5 Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr
 50 55 60
 Lys Pro Arg Glu Glu Gln Phe Asn Ser Thr Tyr Arg Val Val Ser
 65 70 75
 10 Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr
 80 85 90
 Lys Cys Lys Val Ser Asn Lys Gly Leu Pro Ser Ser Ile Glu Lys
 95 100 105
 Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr
 110 115 120
 15 Thr Leu Pro Pro Ser Gln Glu Glu Met Thr Lys Asn Gln Val Ser
 125 130 135
 Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val
 140 145 150
 20 Glu Trp Glx Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr
 155 160 165
 Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Arg
 170 175 180
 Leu Thr Val Asp Lys Ser Arg Trp Gln Glu Gly Asn Val Phe Ser
 185 190 195
 25 Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys
 200 205 210
 Ser Leu Ser Leu Ser Leu Gly Lys
 215 218
 30 <210> 8
 <211> 215
 <212> PRT
 <213> Mus musculus
 <400> 8
 35 Thr Val Pro Glu Val Ser Ser Val Phe Ile Phe Pro Pro Lys Pro
 1 5 10 15
 Lys Asp Val Leu Thr Ile Thr Leu Thr Pro Lys Val Thr Cys Val
 20 25 30
 Val Val Asp Ile Ser Lys Asp Asp Pro Glu Val Gln Phe Ser Trp
 35 40 45
 40 Phe Val Asp Asp Val Glu Val His Thr Ala Gln Thr Gln Pro Arg
 50 55 60
 Glu Glu Gln Phe Asn Ser Thr Phe Arg Ser Val Ser Glu Leu Pro
 65 70 75
 Ile Met His Gln Asp Cys Leu Asn Gly Lys Glu Phe Lys Cys Arg

	80	85	90
	Val Asn Ser Ala Ala Phe Pro Ala Pro	Ile Glu Lys Thr Ile Ser	
	95	100	105
5	Lys Thr Lys Gly Arg Pro Lys Ala Pro	Gln Val Tyr Thr Ile Pro	
	110	115	120
	Pro Pro Lys Glu Gln Met Ala Lys Asp	Lys Val Ser Leu Thr Cys	
	125	130	135
	Met Ile Thr Asp Phe Phe Pro Glu Asp	Ile Thr Val Glu Trp Gln	
	140	145	150
10	Trp Asn Gly Gln Pro Ala Glu Asn Tyr	Lys Asn Thr Gln Pro Ile	
	155	160	165
	Met Asp Thr Asp Gly Ser Tyr Phe Val	Tyr Ser Lys Leu Asn Val	
	170	175	180
15	Gln Lys Ser Asn Trp Glu Ala Gly Asn Thr	Phe Thr Cys Ser Val	
	185	190	195
	Leu His Glu Gly Leu His Asn His His	Thr Glu Lys Ser Leu Ser	
	200	205	210
	His Ser Pro Gly Lys		
	215		
20	<210> 9		
	<211> 218		
	<212> PRT		
	<213> Mus musculus		
	<400> 9		
25	Pro Ala Pro Asn Leu Leu Gly Gly Pro	Ser Val Phe Ile Phe Pro	
	1	5	10
	Pro Lys Ile Lys Asp Val Leu Met Ile	Ser Leu Ser Pro Ile Val	
	20	25	30
30	Thr Cys Val Val Val Asp Val Ser Glu	Asp Asp Pro Asp Val Gln	
	35	40	45
	Ile Ser Trp Phe Val Asn Asn Val Glu	Val His Thr Ala Gln Thr	
	50	55	60
	Gln Thr His Arg Glu Asp Tyr Asn Ser	Thr Leu Arg Val Val Ser	
	65	70	75
35	Ala Leu Pro Ile Gln His Gln Asp Trp	Met Ser Gly Lys Glu Phe	
	80	85	90
	Lys Cys Lys Val Asn Asn Lys Asp Leu	Pro Ala Pro Ile Glu Arg	
	95	100	105
40	Thr Ile Ser Lys Pro Lys Gly Ser Val	Arg Ala Pro Gln Val Tyr	
	110	115	120
	Val Leu Pro Pro Pro Glu Glu Glu Met	Thr Lys Lys Gln Val Thr	
	125	130	135
	Leu Thr Cys Met Val Thr Asp Phe Met	Pro Glu Asp Ile Tyr Val	
	140	145	150

Glu Trp Thr Asn Asn Gly Lys Thr Glu Leu Asn Tyr Lys Asn Thr
 155 160 165
 Glu Pro Val Leu Asp Ser Asp Gly Ser Tyr Phe Met Tyr Ser Lys
 170 175 180
 5 Leu Arg Val Glu Lys Lys Asn Trp Val Glu Arg Asn Ser Tyr Ser
 185 190 195
 Cys Ser Val Val His Glu Gly Leu His Asn His His Thr Thr Lys
 200 205 210
 10 Ser Phe Ser Arg Thr Pro Gly Lys
 215 218
 <210> 10
 <211> 218
 <212> PRT
 <213> Mus musculus
 15 <400> 10
 Pro Ala Pro Asn Leu Glu Gly Gly Pro Ser Val Phe Ile Phe Pro
 1 5 10 15
 Pro Asn Ile Lys Asp Val Leu Met Ile Ser Leu Thr Pro Lys Val
 20 25 30
 20 Thr Cys Val Val Val Asp Val Ser Glu Asp Asp Pro Asp Val Gln
 35 40 45
 Ile Ser Trp Phe Val Asn Asn Val Glu Val His Thr Ala Gln Thr
 50 55 60
 25 Gln Thr His Arg Glu Asp Tyr Asn Ser Thr Ile Arg Val Val Ser
 65 70 75
 His Leu Pro Ile Gln His Gln Asp Trp Met Ser Gly Lys Glu Phe
 80 85 90
 Lys Cys Lys Val Asn Asn Lys Asp Leu Pro Ser Pro Ile Glu Arg
 95 100 105
 30 Thr Ile Ser Lys Pro Lys Gly Leu Val Arg Ala Pro Gln Val Tyr
 110 115 120
 Thr Leu Pro Pro Pro Ala Glu Gln Leu Ser Arg Lys Asp Val Ser
 125 130 135
 35 Leu Thr Cys Leu Val Val Gly Phe Asn Pro Gly Asp Ile Ser Val
 140 145 150
 Glu Trp Thr Ser Asn Gly His Thr Glu Glu Asn Tyr Lys Asp Thr
 155 160 165
 Ala Pro Val Leu Asp Ser Asp Gly Ser Tyr Phe Ile Tyr Ser Lys
 170 175 180
 40 Leu Asn Met Lys Thr Ser Lys Trp Glu Lys Thr Asp Ser Phe Ser
 185 190 195
 Cys Asn Val Arg His Glu Gly Leu Lys Asn Tyr Tyr Leu Lys Lys
 200 205 210
 Thr Ile Ser Arg Ser Pro Gly Lys

215 218

<210> 11
 <211> 218
 <212> PRT
 5 <213> Mus musculus

<400> 11
 Pro Pro Gly Asn Ile Leu Gly Gly Pro Ser Val Phe Ile Phe Pro
 1 5 10 15

10 Pro Lys Pro Lys Asp Ala Leu Met Ile Ser Leu Thr Pro Lys Val
 20 25 30

Thr Cys Val Val Val Asp Val Ser Glu Asp Asp Pro Asp Val His
 35 40 45

Val Ser Trp Phe Val Asp Asn Lys Glu Val His Thr Ala Trp Thr
 50 55 60

15 Gln Pro Arg Glu Ala Gln Tyr Asn Ser Thr Phe Arg Val Val Ser
 65 70 75

Ala Leu Pro Ile Gln His Gln Asp Trp Met Arg Gly Lys Glu Phe
 80 85 90

20 Lys Cys Lys Val Asn Asn Lys Ala Leu Pro Ala Pro Ile Glu Arg
 95 100 105

Thr Ile Ser Lys Pro Lys Gly Arg Ala Gln Thr Pro Gln Val Tyr
 110 115 120

Thr Ile Pro Pro Pro Arg Glu Gln Met Ser Lys Lys Lys Val Ser
 125 130 135

25 Leu Thr Cys Leu Val Thr Asn Phe Phe Ser Glu Ala Ile Ser Val
 140 145 150

Glu Trp Glu Arg Asn Gly Glu Leu Glu Gln Asp Tyr Lys Asn Thr
 155 160 165

30 Pro Pro Ile Leu Asp Ser Asp Gly Thr Tyr Phe Leu Tyr Ser Lys
 170 175 180

Leu Thr Val Asp Thr Asp Ser Trp Leu Gln Gly Glu Ile Phe Thr
 185 190 195

Cys Ser Val Val His Glu Ala Leu His Asn His His Thr Gln Lys
 200 205 210

35 Asn Leu Ser Arg Ser Pro Gly Lys
 215 218

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
20 July 2000 (20.07.2000)

PCT

(10) International Publication Number
WO 00/42072 A3

(51) International Patent Classification⁷: **C07K 16/00**,
A61K 39/395, C12N 15/13, 15/85, 5/10, G01N 33/68,
33/577, C07K 16/22 // 16/28

(21) International Application Number: **PCT/US00/00973**

(22) International Filing Date: 14 January 2000 (14.01.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/116,023 15 January 1999 (15.01.1999) US

(71) Applicant: **GENENTECH, INC.** [US/US]; 1 DNA Way,
South San Francisco, CA 94080-4990 (US).

(72) Inventor: **PRESTA, Leonard, G.**; 1900 Gough Street,
#206, San Francisco, CA 94109 (US).

(74) Agents: **LEE, Wendy, M.** et al.; Genentech, Inc., 1 DNA
Way, South San Francisco, CA 94080-4990 (US).

(81) Designated States (*national*): AE, AL, AM, AT, AU, AZ,
BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK,
DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL,
IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU,
LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT,
RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA,
UG, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent
(AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent
(AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU,
MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM,
GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— With international search report.

(88) Date of publication of the international search report:
30 November 2000

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 00/42072 A3

(54) Title: **POLYPEPTIDE VARIANTS WITH ALTERED EFFECTOR FUNCTION**

(57) Abstract: The present invention concerns polypeptides comprising a variant Fc region. More particularly, the present invention concerns Fc region-containing polypeptides that have altered effector function as a consequence of one or more amino acid modifications in the Fc region thereof.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/00973

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07K16/00 A61K39/395 C12N15/13 C12N15/85 C12N5/10
G01N33/68 G01N33/577 C07K16/22 //C07K16/28

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

BIOSIS, EPO-Internal, PAJ

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5 624 821 A (WINTER GREGORY P ET AL) 29 April 1997 (1997-04-29) column 1, line 44 -column 2, line 61 column 5, line 53-65 column 6, line 38-42 claims --- -/--	1-49



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

6 July 2000

Date of mailing of the international search report

25/07/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040. Tx. 31 651 epo nl.
Fax: (+31-70) 340-3016

Authorized officer

Covone, M

INTERNATIONAL SEARCH REPORT

In: International Application No

PCT/US 00/00973

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>SARMAY G ET AL: "MAPPING AND COMPARISON OF THE INTERACTION SITES ON THE FC REGION OF IGG RESPONSIBLE FOR TRIGGERING ANTIBODY DEPENDENT CELLULAR CYTOTOXICITY ADCC THROUGH DIFFERENT TYPES OF HUMAN FC-GAMMA RECEPTOR"</p> <p>MOLECULAR IMMUNOLOGY, vol. 29, no. 5, 1992, pages 633-639, XP000915240 ISSN: 0161-5890 abstract figure 3 page 634, right-hand column, paragraphs 3,4 page 637, left-hand column, line 15-18 page 638, left-hand column, paragraph 4 -right-hand column, paragraph 2</p> <p>---</p>	1-4, 8-10, 38-42, 44-49
X	<p>WO 98 23289 A (GEN HOSPITAL CORP ;UNIV BRANDEIS (US)) 4 June 1998 (1998-06-04)</p> <p>claims 1-8 example 2</p> <p>---</p>	1-4, 8-15, 31-46
X	<p>GHETIE VICTOR ET AL: "Increasing the serum persistence of an IgG fragment by random mutagenesis."</p> <p>NATURE BIOTECHNOLOGY, vol. 15, no. 7, 1997, pages 637-640, XP000876642 ISSN: 1087-0156 abstract page 637, left-hand column, paragraph 3 page 638, left-hand column, paragraph 1 page 639, right-hand column, paragraph 5 figure 2; table 1</p> <p>---</p>	1-4, 8-15, 31-46
X	<p>MEDESAN CORNELIU ET AL: "Comparative studies of rat IgG to further delineate the Fc:FcRn interaction site."</p> <p>EUROPEAN JOURNAL OF IMMUNOLOGY, vol. 28, no. 7, July 1998 (1998-07), pages 2092-2100, XP000915239 ISSN: 0014-2980 abstract page 2095, right-hand column, paragraph 2 -page 2096, left-hand column, paragraph 1 page 2096, right-hand column, paragraph 2 -page 2097, column 2 page 2098, right-hand column, paragraph 3 page 2098, right-hand column, paragraph 2 page 3</p> <p>---</p> <p>-/--</p>	1-4, 8-15, 31-46

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/00973

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE BIOSIS 'Online! BIOSCIENCES INFORMATION SERVICE, PHILADELPHIA, PA, US; 16 December 1997 (1997-12-16) HENRY ALISTAIR J ET AL: "Participation of the N-terminal region of Cepsilon3 in the binding of human IgE to its high-affinity receptor FepsilonRI." Database accession no. PREV199800070240 XP002142066 abstract & BIOCHEMISTRY, vol. 36, no. 50, 16 December 1997 (1997-12-16), pages 15568-15578, ISSN: 0006-2960</p>	1,2
P,X	<p>WO 99 51642 A (GENENTECH INC) 14 October 1999 (1999-10-14) claims; examples 1-3</p>	1-49

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 00/00973

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 5624821 A	29-04-1997	AT 94171 T	15-09-1993
		AU 600575 B	16-08-1990
		AU 1480388 A	10-10-1988
		DE 3883899 D	14-10-1993
		DE 3883899 T	31-03-1994
		EP 0307434 A	22-03-1989
		EP 0351410 A	24-01-1990
		WO 8807089 A	22-09-1988
		WO 8807054 A	22-09-1988
		GB 2209757 A,B	24-05-1989
		JP 1502875 T	05-10-1989
		US 5648260 A	15-07-1997
WO 9823289 A	04-06-1998	NONE	
WO 9951642 A	14-10-1999	AU 3369999 A	25-10-1999

THIS PAGE BLANK (USPTO)